



# Protective Effect of Food Products Enriched with Unsaponifiable Matter from Palm Fatty Acid Distillate on the Aorta of Hypercholesterolemic Rats

Kgs Ahmadi<sup>1</sup>, Angela Wulansari<sup>2</sup>, Yunianta Subroto<sup>3</sup>, Teti Estiasih<sup>3\*</sup>

<sup>1</sup>Department of Agroindustry Technology, Faculty of Agriculture, Tribhuwana Tunggaladewi University, Jl. Telaga Warna, Tlogomas, Malang, East Java, Indonesia. <sup>2</sup>Master Program of Agricultural Product Technology, Faculty of Agricultural Technology, Brawijaya University, Jl. Veteran, Malang, East Java, Indonesia. <sup>3</sup>Department of Food Science and Technology, Faculty of Agricultural Technology, Brawijaya University, Jl. Veteran, Malang, East Java, Indonesia.

---

## ARTICLE INFO

### Article history:

Received on: 12/08/2017

Accepted on: 25/09/2017

Available online: 30/12/2017

### Key words:

Palm fatty acid distillate, unsaponifiable matter, enrichment, hypercholesterolemic, atherosclerosis.

---

## ABSTRACT

Palm oil refining process produced palm fatty acid distillate (PFAD) as a by-product in deodorization stage. Saponification of PFAD produced unsaponifiable matter (USM) which was rich in vitamin E mainly tocotrienols, phytosterols, and squalene. This study evaluated the therapeutic effect of food products (instant noodle, bread, and biscuit) enriched with bioactive compounds from USM on the aorta of hypercholesterolemic rats. Rats were fed with atherogenic diet for 14 days to have blood total cholesterol  $\geq 200$  mg/dl. Rats then were fed according to each treatment group for 8 weeks. Total cholesterol, HDL (high density lipoprotein) cholesterol, and LDL (low density lipoprotein) cholesterol were analyzed at the end of experiment. Rats were then sacrificed and aorta abdominal was collected for histopathological study. The result showed total cholesterol and LDL cholesterol of rats fed by USM enriched food products were lower than that of corresponding products. Also, the ratio of total cholesterol to HDL and LDL cholesterol to HDL were better. Rats fed with USM enriched food products had a better aortic histopathological image than rats fed with non-enriched food products. Rats fed with USM enriched foods had less severe morphological lesions of the aortic wall with less foam cells in tunica intima, less fat deposits in tunica media, elongated nuclei and organized myofibrils. This study indicated bioactive compounds in food products enriched with USM of PFAD offered good therapeutic effect against atherosclerosis development of hypercholesterolemic rats. USM enriched biscuit revealed the best therapeutic effect.

---

## INTRODUCTION

Palm oil contains minor components such as carotenoids, vitamin E (tocopherol and tocotrienols), phytosterols, squalene, and phenolic compounds (Mukherjee and Mitra, 2009; Loganathan *et al.*, 2009; Che *et al.*, 2014). Deodorization stage on palm oil refining process is aimed to remove undesirable odor and free fatty acid, which also produces palm fatty acid distillate (PFAD) as a by-product. PFAD contains

free fatty acids and other minor components. Estiasih *et al.* (2013) reported PFAD was consisted of 85-90% fatty acids as main compound and bioactive compounds such as vitamin E (60-200 ppm), phytosterols (400-7500 ppm), and squalene (400-2800 ppm) as minor components. Separation of free fatty acids and bioactive compounds in PFAD can be occurred by saponification reaction. Fatty acids are hydrolyzed by the base to produce soap and other unsaponifiable components are dissolved in unsaponifiable fraction (Fernandes and Cabral, 2007). The study by Estiasih *et al.* (2014) showed unsaponifiable matters of PFAD was contained vitamin E 19600 ppm, phytosterols 5500 ppm, and squalene 323000 ppm.

---

\* Corresponding Author

Email: [teties@yahoo.co.id](mailto:teties@yahoo.co.id)

Research by Ahmadi *et al.* (2012) showed unsaponifiable matters of PFAD was contained 35000 ppm vitamin E that consisted of 11844 ppm  $\alpha$ -tocopherol, 7441 ppm  $\alpha$ -tocotrienol, 2250.5 ppm  $\delta$ -tocotrienol, and 13464.5 ppm  $\gamma$ -tocotrienol.

Vitamin E, phytosterols, and squalene from USM of PFAD have a potential hypocholesterolemic effect. Estiasih *et al.* (2014) reported USM of PFAD decreased total cholesterol, triglyceride, LDL cholesterol, and increased HDL cholesterol level in hypercholesterolemic rats. Cholesterol, especially LDL cholesterol is a major risk factor for atherosclerosis. The presence of HDL cholesterol will affect the atherogenic degree of LDL cholesterol, therefore atherosclerosis could be prevented by improving lipid profiles (Tomkin and Owens, 2012).

Vitamin E in USM of PFAD is dominated by tocotrienols. Study by Qureshi *et al.* (2002) showed tocotrienols rich fraction supplementation decreased total cholesterol, LDL cholesterol, and triglyceride levels in human subjects with hypercholesterolemia. Oxidized LDL is the major cause of atherosclerosis. Vitamin E as antioxidant prevented oxidation of LDL cholesterol in *in vitro* study (Niki, 2011). A study in hyperlipidemic rats showed the group of rats treated with vitamin E had smaller atherosclerotic area than groups without vitamin E treatment (Hasty *et al.*, 2007).

Phytosterols have the ability to compete with cholesterol during the formation of micelles in the intestine. Rats that given additional phytosterols in high-fat diet had lower total cholesterol, LDL cholesterol, and triglyceride levels (Awaisheh *et al.*, 2013). Squalene, as an antioxidant, prevents free radical damage in cells (Buddhan *et al.*, 2007). Squalene also reported to decrease cholesterol levels in plasma (Bombo *et al.*, 2013), inhibit atherosclerosis formation, and decrease cholesterol levels in liver (Guillen *et al.*, 2008).

Increasing number of chronic diseases such as cardiovascular disease, diabetes mellitus, and cancer, leads to increase promotion of enriched foods that have a health benefit in form of functional foods (Jew *et al.*, 2009). Fortification of phytosterols have been done in some food products such as bread, cereals, and margarine which were decreased LDL cholesterol in human subject with hypercholesterolemia (Nestel *et al.*, 2001). Fortification of vitamin E in milk (Hayes *et al.*, 2001) and margarine (van het Hof *et al.*, 1998) prevented LDL cholesterol oxidation and increased LDL cholesterol resistance to oxidation. The addition of high squalene containing food such as amaranth in pasta (Martinez *et al.*, 2016), and rice bran oil in biscuit (Bhanger *et al.*, 2008) increased squalene content of the food products and made them as a good source of antioxidant. USM of PFAD that contained phytosterols, squalene, and vitamin E is possible to use as ingredient for enriching foods to have hypocholesterolemic effect and aortic protection.

This study examined therapeutic effect of USM enriched instant noodle, bread, and biscuit against atherosclerosis development in hypercholesterolemic rats.

## MATERIALS AND METHOD

### Materials

PFAD kindly obtained from PT Salim Ivonas Pratama, Surabaya, Indonesia. Reagents that used in this study were the technical grade reagents for preparation of USM and other chemical reagents for preparation of fresh specimens. Cholesterol kit (CHOD PAP) was used for lipid profile analysis. Other materials were instant noodle, bread, and biscuit ingredients, standard diet AIN-93M, and atherogenic diet ingredients (cholesterol, cholic acids, and tallow).

### Preparation of USM from PFAD

Preparation and saponification of PFAD was according to the method of Estiasih *et al.* (2014).

### Preparation of USM enriched Food Products

Instant noodle was made by mixing wheat flour, eggs, salt, water, and alkaline water until smooth. USM of PFAD as much as 1% of the dough (w/w) was added and mixed well. The dough was rested for 20 min then sheeted and slotted until had the strands of noodle. Noodle strands were steamed for 10 min at 100°C and fried in palm oil for 15 s.

Bread was made by mixing dry ingredients such as wheat flour, sugar, milk powder and instant yeast. Egg yolks, margarine, and warm water were added to the dry ingredients and mixed. USM as much as 1% weight of dough (w/w) was added to the dough and mixed until smooth. The dough then rested for 30 min. The dough was divided into 100 g and baked for 20 min at 180°C.

Biscuit was prepared by mixing butter, margarine, egg yolk and sugar powder for 3 mins. Wheat flour, baking powder, salt and vanilla then added and mixed until smooth. USM as much as 1% of the dough (w/w) added and mixed well. The dough was sheeted and cut into 4x4x1 cm, then baked for 15 min at 180°C.

All food products were analyzed for bioactive compounds including vitamin E, phytosterols, and squalene levels.

### Animals Preparation and Treatment

This study was approved for Ethical Clearance No. 608-KEP-UB from Animal and Care and Use Committee, Brawijaya University. About 40 male Wistar rats, age 2-3 months, weight 150-200 g, were adapted for 1 week and fed ad libitum with standard diet AIN-93M. After the adaptation period, rats were divided into 8 groups consisted of one normal group and seven hypercholesterolemic groups. Groups of hypercholesterolemic rats were given atherogenic diet whereas the group of normal rats was given standard diet AIN-93M (Table 1). Atherogenic diet was given for 14 days. After rats from hypercholesterolemic groups had reached blood total cholesterol level  $\geq 200$  mg/dl, atherogenic diet was stopped and all groups of rats were fed according to each treatment group for 8 weeks. At week 8, blood serum total cholesterol, HDL cholesterol, and LDL cholesterol of rats was analyzed photometrically by CHOD-PAP method (Cholesterol

Oxidase-Phenol Aminophenazone) (Siedel *et al.*, 1983). The ratio of total cholesterol/HDL cholesterol and LDL cholesterol/HDL cholesterol were analyzed as the indicator of risk factors of coronary heart disease.

Treatment groups consisted of normal group of rats fed with standard diet AIN-93M (K1), hypercholesterolemic rats fed with non enriched instant noodle (K2), hypercholesterolemic rats fed with non enriched bread (K3), hypercholesterolemic rats fed with non enriched biscuit (K4), hypercholesterolemic rats fed with USM enriched instant noodle (K5), hypercholesterolemic rats fed with USM enriched bread (K6), hypercholesterolemic rats fed with USM enriched biscuit (K7), and hypercholesterolemic rats fed with standard diet AIN-93M (K8). In the end of week-8, all rats were sacrificed and abdominal aorta from the heart to the liver was collected.

**Table 1:** Composition of standard diet AIN-93M and atherogenic diet.

Ingredient	Composition (g/kg Diet)	
	AIN-93M	Atherogenic Diet
Corn starch	620.692	620.692
casein	140	116.528
Sucrose	100	100
Soy bean oil	40	40
CMC	50	50
Mineral mix	35	35
Vitamin mix	10	10
L-cystin	1.8	1.8
Choline bitartrate	2.5	2.5
TBHQ	-	-
Cholic Acid	-	2
Cholesterol	-	20
Tallow	-	185

Source: Reeves *et al.* (1993).

### Aorta Histopathology Analysis

Aorta samples were prepared according to the method of Thent *et al.* (2012). Embedding samples were performed on paraffin wax with tissue embedding center (Tissue-Tek TEC 5, Sakura, Japan). Aorta was cut in 5  $\mu$ m thickness of with 3 replications for each rats using microtome (LEICA RM2245). Samples were stained with Haematoxylin and Eosin (H&E) (Tissue-Tek DRS 2000, Sakura, Japan) and then analyzed under the microscope (Olympus XC 10, Tokyo, Japan) with 400x enlargement. The histopathology image was analyzed using software OlyVIA 2.2. Intima media thicknesses (IMT) was measured with dotSlide virtual microscopy system in four different section (0°, 90°, 180°, 270°).

### Analysis of Bioactive Compounds

Phytosterols analysis was followed method of Khatoon *et al.* (2010), vitamin E analysis was according to Ball (1988), and squalene analysis was using GC MS (GCMS-QP2010S Shimadzu) with derivatization method according to Mendez *et al.* (2003).

### Data Analysis

Data were analyzed for variance test (ANOVA) and further tested by Duncan Multiple Range Test (DMRT) using software SPSS.

## RESULT AND DISCUSSION

### Bioactive Compounds

Table 2 showed all food products contained bioactive compounds from USM. Predominant bioactive compound in USM was phytosterols. The level of squalene was higher than vitamin E in USM (data not shown). Among food products there was different bioactive content. The highest amount of phytosterols and squalene was found in biscuit. Meanwhile, the highest quantity of vitamin E was in instant noodle. Bread showed the lowest bioactive compounds among all foods. All vitamin E in all food products is tocotrienols and no tocopherols was detected.

**Table 2:** Bioactive compounds of USM enriched food products.

Bioactive Compound (mg/100 g)	Product		
	Biscuit	Instant Noodle	Bread
Total vitamin E (tocotrienols)	23.46±0.71	35.76±1.09	22.68±0.05
Total phytosterols	261.35±5.01	186.43±3.71	122.11±2.59
Squalene	128.45±2.26	72.51±1.42	56.29±1.25

### Total Cholesterol Level

Table 3 showed blood serum total cholesterol level after 8 week feeding. The lowest total cholesterol was found in normal group and the highest was hypercholesterolemia group, both were fed by standard diet. Group of rats treated by USM enriched products showed lower total cholesterol level than their corresponding food. It means that USM enriched food products decreased total cholesterol. The highest decrease was found in USM enriched bread compared to non-enriched. Compared to hypercholesterolemia group fed by standard diet, all hypercholesterolemia group fed by non-enriched biscuit, bread, and instant noodle showed lower total cholesterol level.

Ratio of total cholesterol to HDL cholesterol and LDL cholesterol to HDL cholesterol were indicators of the risk of coronary vascular diseases (CVD) (Fernandez and Webb, 2008). Hypercholesterolemic group which fed by standard diet showed the highest ratio, meanwhile the normal group fed by standard diet revealed the lowest. All food products caused decrease in the ratio although without enrichment of USM. Compared to non-enriched foods, all enriched foods showed better ratio of total cholesterol to HDL cholesterol and LDL cholesterol to HDL cholesterol. This is an indicator that enrichment of foods by USM improved lipid profile and lower the risk of CVD. Among all USM enriched foods, bread had the best ratio although the level of bioactive compounds in bread was the lowest.

### Aorta Histopathology

Atherogenic diet was given for 14 days to raise blood cholesterol levels in hypercholesterolemic groups. Beside raising blood cholesterol levels, atherogenic diet also affected aortic histopathology. The normal group of rats (K1) showed a normal shape of the aorta with none foam cells, none visible fat accumulation or vacuolization in intima media layer, and none development of aortic plaque (Figure 1).



**Fig. 1:** Aortic histopathological image of normal group of rats + AIN-93M (K1) showing normal tunica intima (TI), tunica media (TM), and tunica adventitia (TA) (enlargement: 400x, scale bar: 20 $\mu$ m).

A group of hypercholesterolemic rats fed with standard diet (K8) showed a different result. There were irregular tunica intima with a development of plaque and foam cells. Some fat deposits and dense irregular nuclei were found in tunica media. The elastic fibers were wavy and interrupted (Figure 8). Atherosclerosis is caused by elevated blood cholesterol levels especially low-density lipoprotein (LDL) cholesterol and the appearance of free radicals; reactive oxygen species (ROS); reactive nitrogen species (RNS) (Niki, 2011).

This group (K8) also revealed high level of LDL cholesterol and low level of HDL cholesterol (Table 3). Also, the ratio of total cholesterol to HDL cholesterol and LDL cholesterol to HDL cholesterol was the highest. Both ratio was an indicator of the risk of coronary vascular disease that usually initiated by aortic plaque formation. High level of total cholesterol, LDL cholesterol, and low level of HDL cholesterol were probably as the cause of plaque development in aorta of this group.

Rats fed with USM enriched food products showed better aortic histopathological image than non-enriched foods. USM enriched food groups had less fat deposit in tunica media, elastic fibers were straight and less wavy, foam cell also only found in USM enriched bread group (Figure 5, 6, 7). While non-enriched foods groups showed more fat deposit in tunica media, more foam cells in tunica intima, elastic fibers were wavy and slightly interrupted (Figure 2, 3, 4). Rats fed with USM enriched food products had less severe morphological lesion and the aorta looked similar with normal group.


Profile lipid has main role in atherosclerosis development. In the end of week 8, profile lipid of rats fed with USM enriched food products was better than that of non-enriched food products. Nevertheless, rats fed with USM enriched food products had better profile lipid than non-enriched food products, as well as lower ratio of total cholesterol to HDL cholesterol and ratio of LDL cholesterol to HDL cholesterol (Table 3). Better aortic histopathology possibly was related to better profile lipid improvement in rats fed with USM enriched food products. The best ratio of total cholesterol to HDL cholesterol and LDL cholesterol to HDL cholesterol was found in group of rats fed by USM enriched bread. However, among USM enriched foods, the histopathological analysis revealed that the best therapeutic effect was found in USM enriched biscuit and the worst was in USM enriched bread. It seemed that in this case, the ratio of total cholesterol to HDL cholesterol and LDL cholesterol to HDL cholesterol not the only indicator for the aortic plaque development. Food intake also affected the therapeutic effect of USM enriched foods. Among USM enriched foods, intake of bread was the lowest (data not shown) that meant the intake of the bioactive compounds from bread also the lowest. Moreover, bread also showed the lowest fat content (data not shown) among USM enriched food products. Presumably, less fat intake resulted in better lipid profile and cholesterol ratio.

Previous research reported USM of PFAD contained bioactive compounds such as vitamin E (tocopherol and tocotrienols), phytosterols, and squalene (Estiasih *et al.*, 2014; Ahmadi *et al.*, 2012). Table 2 showed that all USM enriched food contained tocotrienols, phytosterols, and squalene. Bioactive compounds in USM presumed had a protective effect on aorta wall from radical damage and inhibited lipid oxidation. Aortic histopathological image of rats fed with USM enriched biscuit (K7) and instant noodle (K5) were slightly better than rats fed with USM enriched bread (K6) (Figure 5, 6, 7). USM enriched biscuit had the highest bioactive compounds and USM enriched bread had the lowest bioactive compounds (Table 2). The amount of bioactive compounds was assumed to be the cause of different therapeutic effect among USM enriched foods against atherosclerosis development. However, the existence of bioactive compounds in USM enriched food products, surely gave significant impact on prevention of atherosclerosis development compare with non-enriched food products.


**Table 3:** Total cholesterol level of group of rats after 8-week feeding.

Group	Total Cholesterol (mg/dl)	LDL Cholesterol (mg/dl)	HDL Cholesterol (mg/dl)	Total Cholesterol/HDL Cholesterol	LDL Cholesterol/HDL Cholesterol
K1 = Normal Group, Standard Diet	95.21 $\pm$ 3.17 a	29.89 $\pm$ 3.50 a	79.56 $\pm$ 2.51 f	1.20 $\pm$ 0.00 h	0.37 $\pm$ 0.03 g
K2 = Hypercholesterolemia Group, Instant Noodle	142.11 $\pm$ 8.05 e	44.66 $\pm$ 2.01 d	53.20 $\pm$ 3.62 b	2.67 $\pm$ 0.03 b	0.84 $\pm$ 0.02 b
K3 = Hypercholesterolemia Group, Bread	125.78 $\pm$ 6.69 d	37.09 $\pm$ 1.43 bc	60.44 $\pm$ 2.62 c	2.08 $\pm$ 0.02 e	0.61 $\pm$ 0.00 d
K4 = Hypercholesterolemia Group, Biscuit	126.61 $\pm$ 3.76 d	39.30 $\pm$ 1.65 bc	55.00 $\pm$ 2.51 b	2.30 $\pm$ 0.04 d	0.71 $\pm$ 0.00 c
K5 = Hypercholesterolemia Group, USM Enriched Instant Noodle	127.77 $\pm$ 4.94 d	38.33 $\pm$ 1.80 bc	53.23 $\pm$ 1.76 b	2.40 $\pm$ 0.01 c	0.72 $\pm$ 0.01 c
K6 = Hypercholesterolemia Group, USM Enriched Bread	102.31 $\pm$ 4.02 b	31.83 $\pm$ 1.89 a	73.82 $\pm$ 1.69 e	1.39 $\pm$ 0.02 g	0.43 $\pm$ 0.02 f
K7 = Hypercholesterolemia Group, USM Enriched Biscuit	108.93 $\pm$ 4.46 c	36.12 $\pm$ 1.33 b	68.67 $\pm$ 1.98 d	1.63 $\pm$ 0.07 f	0.53 $\pm$ 0.00 e
K8 = Hypercholesterolemia Group, Standard Diet	202.98 $\pm$ 1.99 f	77.64 $\pm$ 2.21 e	25.00 $\pm$ 2.32 a	8.18 $\pm$ 0.68 a	3.12 $\pm$ 0.20 a







**Fig. 2:** Aortic histopathological image of hypercholesterolemic rats + instant noodle (K2) revealing irregular tunica intimae (TI) with foam cells (red arrows), irregular tunica media (TM) with fat deposits (stars) and dense irregular nuclei (black arrow), (enlargement: 400x, scale bar: 20µm).



**Fig. 3:** Aortic histopathological image of hypercholesterolemic rats + bread (K3) showing irregular tunica intimae (TI) with more foam cells (red arrows), irregular tunica media (TM) with more fat deposits (stars), and dense irregular nuclei (black arrows) (enlargement: 400x, scale bar: 20µm).




**Fig. 4:** Aortic histopathological image of hypercholesterolemic rats + biscuit (K4) revealing irregular tunica intimae (TI) with foam cells (red arrows), irregular tunica media (TM) with fat deposits (stars) and dense irregular nuclei (black arrows) (enlargement: 400x, scale bar: 20µm).




**Fig. 5:** Aortic histopathological image of hypercholesterolemic rats + USM enriched instant noodle (K5) illustrating straight tunica intimae (TI), less fat


deposits in tunica media (TM), and elongated nuclei (black arrow) (enlargement: 400x, scale bar: 20µm).



**Fig. 6:** Aortic histopathological image of hypercholesterolemic rats + USM enriched bread (K6) showing less severe morphological lesions of aortic wall, less foam cells (red arrow), less fat deposits (stars) in tunica media (TM), and elongated nuclei (black arrows) (enlargement: 400x, scale bar: 20µm).



**Fig. 7:** Aortic histopathological image of hypercholesterolemic rats + USM enriched biscuit (K7) revealing straight tunica intimae (TI), less fat deposits (stars) in tunica media (TM), and elongated nuclei (black arrow) (enlargement: 400x, scale bar: 20µm).



**Fig. 8:** Aortic histopathological image of hypercholesterolemic rats + standard diet AIN-93M (K8) showing irregular tunica intimae (TI) with foam cells and plaque development (red arrows), irregular tunica media (TM) with fat deposits (stars) and dense irregular nuclei (black arrows), (enlargement: 400x, scale bar: 20µm).

Vitamin E as antioxidant has the ability to decrease the level of oxidized LDL and expression of CD38 which had a role in foam cell formation (Tang *et al.*, 2014). The present study showed rats fed with USM enriched food products had none foam cells except group fed with USM enriched bread, which contained the lowest amount of vitamin E. Vitamin E also can decrease homocysteine in the blood and prevent the proliferation of smooth muscle cells (SMC) (Kirac *et al.*, 2013). High level of

homocysteine and cholesterol in blood will increase proliferation of SMC and oxidative stress which induced atherosclerosis (Ozer *et al.*, 2003). Supplementation of vitamin E in rats increased expression of peroxisome proliferator activated receptor gamma (PPAR $\gamma$ ), liver-x-receptor- $\alpha$  (LXR $\alpha$ ), and ATP-binding cassette transporter ABCA1. Increased PPAR $\gamma$  gene expression stimulated LXR $\alpha$  activation. PPAR $\gamma$  and LXR $\alpha$  stimulated more of ABCA1 gene expression. ABCA1 removed cholesterol from macrophage and preventing the appearance of foam cell (Tang *et al.*, 2014). In this study, tocotrienols were the most responsible in preventing LDL oxidation since no tocopherols was detected in all food products. The addition of USM to instant noodle, bread, and biscuit also enriched the food products with phytosterols and squalene. Squalene as the antioxidant, protected cell membrane lipid bilayer from free radical damage (Farvin *et al.*, 2007). Rats fed with high cholesterol diet and phytosterols had lower blood serum cholesterol levels, less damage in arterial walls, lower lipid accumulation in arterial walls, and less macrophages appearance on the arterial walls compared to control group (Bombo *et al.*, 2013). The human study had shown that phytosterols increased circulation of endothelial progenitor cells (EPCs) that have a function for regeneration and protection of endothelial cells to prevent atherosclerosis (Chen *et al.*, 2015). In this study, groups of rats fed with USM enriched foods had less fat deposits in the aortic wall and less endothelial lesion.


Vitamin E, phytosterols, and squalene as bioactive compounds in USM inhibited atherosclerosis development. Aortic histopathology image of hypercholesterolemic rats fed with USM enriched food products was the proof of its therapeutic effect. Hypercholesterolemic rats fed with non-enriched food products showed less improvement in histopathological alterations. The damage of aortic in these groups was almost same with hypercholesterolemic rats those were fed by the standard diet. In the other side, hypercholesterolemic rats fed with USM enriched food product had less aortic damage and almost had a similar histopathological image with normal rats.

### Intimae Media Thickness (IMT)

The present study revealed atherogenic diet treatment for 14 days increased blood total cholesterol levels but not sufficient to induce thickening of aortic wall. Some groups of hypercholesterolemic rats had lower average of IMT than normal group (Figure 9). Hypercholesterolemic rats fed with USM enriched biscuit had the highest IMT of all, but histopathological image showed this group had less aortic damage than other hypercholesterolemic groups. Average IMT of the normal group also did not significantly different with hypercholesterolemic rats fed the standard diet. Nevertheless, the histopathological image of hypercholesterolemic rats fed with standard diet showed aortic damage and started the atherosclerosis development.

Variation of intimae media thickness is caused by myoelectric (smooth muscle cells) structure and density of aortic layers with 18% degree of variation (Davis *et al.*, 2010).

Atherogenic diet treatment for a longer time may need to induce aortic wall thickening in hypercholesterolemic rats.



**Fig. 9:** Aorta intimae media thickness (IMT) at week 8. K1=normal group, standard diet AIN-93M; K2=hypercholesterolemia, instant noodle; K3=hypercholesterolemic, bread; K4=hypercholesterolemic, biscuit; K5=hypercholesterolemic, USM enriched instant noodle; K6=hypercholesterolemic, USM enriched bread; K7=hypercholesterolemic, USM enriched biscuit, K8=hypercholesterolemic, standard diet AIN-93M.

### CONCLUSION

USM enriched instant noodle, bread, and biscuit inhibited atherosclerosis plaque development in hypercholesterolemic rats. Bioactive compounds in USM of PFAD decreased foam cells development, decreased fat deposits, and improved morphological image of the aortic wall in hypercholesterolemic rats. Groups of hypercholesterolemic rats fed with USM enriched food products had a better aortic histopathological image than non-enriched food products. Between the USM enriched foods group, rats fed with USM enriched biscuit and instant noodle had better histopathological image than USM enriched bread. The best therapeutic effect was found in group of rats fed with USM enriched biscuit. However, overall rats fed with USM enriched food products had less severe aortic lesion than non-enriched foods and almost had a similar aortic histopathological image with the normal group of rats.

Treatment with atherogenic diet for 14 days could not induced atherosclerosis plaque in hypercholesterolemic groups of rats. The aortic IMT was not significantly different between the normal group of rats with the hypercholesterolemic group of rats. Longer time of atherogenic diet feeding may need to induce plaque development.

### ACKNOLEGMENT

**Financial support and sponsorship:** The authors would like to thank and appreciate Directorate of Research and Community Service, Directorate General of Strengthening Research and Development, Ministry of Research, Technology, and Higher Education, for funding this research through Penelitian Strategis Nasional with Contract No. 007/SP2H/LT/DRPM/II/2016, February 17, 2016.

**Conflict of Interests:** There are no conflicts of interest.

## REFERENCES

- Ahmadi K, Kumalaningsih S, Wijana S, Santoso I. Optimizing vitamin E purification from unsaponifiable matter of palm fatty acid distillate by low temperature solvent crystallization. *Journal of Food Science and Engineering*, 2012; 2: 557-563.
- Awaisheh SS, Khalifeh MS, Al-Ruwaili MA, Khalil OM, Al-Ameri OH, Al-Groom R. Effect of supplementation of probiotics and phytosterols alone or in combination on serum and hepatic lipid profiles and thyroid hormones of hypercholesterolemic Rats. *Journal of Dairy Science*, 2013; 96: 9-15.
- Ball GFM. *Fat-Soluble Vitamin Assays in Food Analysis (A Comprehensive Review)*. Elsevier Applied Science, 1988, London.
- Bhanger MI, Iqbal S, Anwar F, Imran M, Akhtar M, Zia-ul-Haq M. Antioxidant potential of rice bran extracts and its effects on stabilisation of cookies under ambient storage. *International of Food Science & Technology*, 2008; 43: 779-786.
- Bombo RP, Afonso MS, Machado RM, Lavrador MS, Nunes VS, Quintao ER, Koike M, Catanozi S, Lin CJ, Nakandakare ER, Lottenberg AM. Dietary phytosterol does not accumulate in the arterial wall and prevents atherosclerosis of LDLr-KO mice. *Atherosclerosis*, 2013; 231(2): 442-447.
- Buddhan S, Sivakumar R, Dhandapani N, Ganesan B, Anandan R. Protective effect of dietary squalene supplementation on mitochondrial function in liver of aged rats. *Prostaglandins Leukot Essent Fatty Acids*, 2007; 76(6): 349-355.
- Che ICA, Karupaiah T, Sundram K, Tan YA, Balasundram N, Leow SS, Nasruddin NS, Sambanthamurthi R. Oil palm phenolics and vitamin E reduce atherosclerosis in rabbits. *Journal of Functional Foods*, 2014; 7(1): 541-550.
- Chen D, Huang P, Chiang C, Leu H, Chen J, Lin S. Phytosterol increase circulating endothelial progenitor cells and insulin-like growth factor-1 levels in patients with nonalcoholic fatty liver disease: A randomized crossover study. *Journal of Functional Foods*, 2015; 13:148-157.
- Davis PH, Dawson JD, Blecha MB, Mastbergen RK, Sonka M. Measurement of aortic intimal-medial thickness in adolescents and young adults. *Ultrasound Med Biol*, 2010; 36(4): 560-565.
- Estiasih T, Ahmadi K, Widyaningsih TD, Maligan JM, Mubarak AZ, Zubaidah E, Mukhlisiyyah J, Puspitasari R. Bioactive compounds of palm fatty acid distillate (PFAD) from several palm oil refineries. *Journal of Food Science and Technology*, 2013; 5(9): 1153-59.
- Estiasih T, Ahmadi K, Widyaningsih TD, Rhitmayanti E, Fidyasari A, Purnomo K, Wahyuni Y. The effect of unsaponifiable fraction from palm fatty acid distillate on lipid profile of hypercholesterolemic rats. *Journal of Food and Nutrition Research*, 2014; 2(12): 1029-1036.
- Farvin, KHS, Anandan R, Kumar SHS, Shiny KS, Mathew S, Sankar TV, Nair PGV. Cardio protective effect of squalene on lipid profile in isoprenaline-induced myocardial infarction in rats. *J.Med.Food*, 2006; 9(4): 531-536.
- Fernandez ML, West KL. Mechanisms by which dietary fatty acids modulate plasma lipids. *J Nutr*, 2005; 135(9): 2075-2078.
- Fernandez ML, Webb D. The LDL to HDL Cholesterol Ratio as a Valuable Tool to Evaluate Coronary Heart Disease Risk. *J Am Coll Nutr*, 2008, 27(1): 1-5.
- Guillen N, Acin S, Navarro MA, Perona JS, Arbones-Mainar JM, Arnal C, Sarria AJ, Surra JC, Carnicer R, Orman I, Segovia JC, Ruiz-Gutierrez V, Osada J. Squalene in a sex-dependent manner modulates atherosclerotic lesion which correlates with hepatic fat content in apoE-knockout male mice. *Atherosclerosis*, 2008; 197(1): 72-83.
- Hasty AH, Gruen ML, Terry ES, Surmi BK, Atkinson RD, Gao L, Morrow JD. Effects of vitamin E on oxidative stress and atherosclerosis in an obese hyperlipidemic mouse model. *J Nutr Biochem*, 2007; 18(2): 127-133.
- Hayes K, Pronczuk A, Perlman D. Vitamin E in enriched cow milk uniquely enriches human plasma lipoproteins. *Am J Clin Nutr*, 2001; 74(2): 211-218.
- Jew S, AbuMweis SS, Jones PJ. Evolution of the human diet: linking our ancestral diet to modern functional foods as a means of chronic disease prevention. *J Med Food*, 2009; 12(5): 925-934.
- Khatoun S, Rajan, RGR, Krishna AGG. Physicochemical characteristics and composition of Indian soybean oil deodorizer distillate and the recovery of phytosterols. *J. Am Oil Chem Soc*, 2010, 87(3): 321-326.
- Kirac D, Negis Y, Ozer NK. Vitamin E attenuates homocystein and cholesterol induced damage in rat aorta. *Cardiovasc Pathol*, 2013; 22(6): 465-472.
- Loganathan R, Selvaduray KR, Radhakrishnan A, Nesaretnam K. Palm oil: rich in health promoting phytonutrients. *Palm Oil Developments*, 2009; 50: 16-25.
- Martinez CS, Ribotta PD, Anon MC, Leon AE. Effect of amaranth flour (*Amaranthus mantegazzianus*) on the technological and sensory quality of bread wheat pasta. *Food Sci Technol Int*, 2014; 20(2):127-135.
- Mendez E, Blanco M, Laguna A, Garcia E. Isolation and characterization of a mixture of higher primary aliphatic alcohols of high molecular weight from henequen (*Agave furcroydes L.*) wax. *Revista CENIC Ciencias Químicas*, 2003, 34(1): 35-38.
- Mukherjee S, Mitra A. Health effects of palm oil. *J Hum Ecol*, 2009; 26(3): 197-203.
- Nestel P, Cehun M, Pomeroy S, Abbey M, Weldon G. Cholesterol-lowering effects of plant sterol esters and non-esterified stanols in margarine, butter and low-fat foods. *Eur J Clin Nutr*, 2001; 55(12): 1084-1090.
- Niki E. Do free radicals play casual role in atherosclerosis? Low density lipoprotein oxidation and vitamin E revisited. *J.Clin Biochem Nutr*, 2011; 48(1): 3-7.
- Ozer NK, Negis Y, Aytan N. Molecular mechanisms of cholesterol or homocysteine effect in the development of atherosclerosis: Role of vitamin E. *BioFactors*, 2003; 19(1-2): 63-70.
- Qureshi AA, Sami SA, Salser WA, Khan FA. Dose-dependent suppression of serum cholesterol by tocotrienol-rich fraction (TRF25) of rice bran in hypercholesterolemic humans. *Atherosclerosis*, 2002; 161(2): 199-207.
- Reeves PG, Nielson FH, Fahey GC Jr. AIN-93 purified diets for laboratory rodents: final report of The American Institute of Nutrition ad hoc Writing Committee on reformulation of the AIN-76A rodent diet. *J Nutr*, 1993; 123(11): 1939-1951.
- Siedel J, Hagele EO, Ziegenhorn J, Wahiefeld AW. Reagent for the enzymatic determination of serum total cholesterol with improved lipolytic efficiency. *Clin Chem*, 1983, 29:1075-1080.
- Tang F, Lu M, Zhang S, Mei M, Wang T, Liu P, Wang H. Vitamin E conditionally inhibits atherosclerosis in ApoE knockout mice by anti-oxidation and regulation of vasculature gene expressions. *Lipids*, 2014; 49(12): 1215-1223.
- Thent ZC, Lin TS, Das S, Zakaria Z. Histological changes in the heart and the proximal aorta in experimental diabetic rats fed with piper sarmentsoum. *Afr J Tradit Complement Altern Med*, 2012; 9(3): 396-404.
- Tomkin GH, Owen D. LDL as a cause of atherosclerosis. *The Open Atherosclerosis & Trombosis Journal*, 2012; 5: 13-21.
- van het Hof KH, Tijburg LB, de Boer HS, Wiseman SA, Weststrate JA. Antioxidant enriched margarine increases the antioxidant status. *Eur J Clin Nutr*, 1998; 52(4): 292-299.

### How to cite this article:

Ahmadi K, Wulansari A, Subroto Y, Estiasih T. Protective Effect of Food Products Enriched with Unsaponifiable Matter from Palm Fatty Distillate on The Aorta of Hypercholesterolemic Rats. *J App Pharm Sci*, 2017; 7 (12): 090-096.