

Triterpenes and Sterol from *Artocarpus ovatus*

Consolacion Y. Ragasa^{1,2}, Jocelyn L. Caro², Chien-Chang Shen³

¹Chemistry Department, De La Salle University Science & Technology Complex Leandro V. Locsin Campus, Biñan City, Laguna, Philippines.

²Chemistry Department De La Salle University, 2401 Taft Avenue, Manila, 1004, Philippines.

³National Research Institute of Chinese Medicine, 155-1, Li-Nong St., Sec. 2, Taipei 112, Taiwan.

ARTICLE INFO

Article history:

Received on: 12/08/2014

Revised on: 26/08/2014

Accepted on: 18/09/2014

Available online: 30/10/2014

Key words:

Artocarpus ovatus, Moraceae, triterpenes, sterol

ABSTRACT

Chemical investigation of the dichloromethane extract of the twigs of *Artocarpus ovatus* afforded lupeol (**1a**), α -amyrin (**1b**), β -amyrin (**1c**), lupeol fatty acid ester (**2a**), α -amyrin fatty acid ester (**2b**), β -myrin fatty acid ester (**2c**), betulin (**3a**), 3β , 28-Dihydroxyolean-12-ene (**3b**), oleanolic acid (**4**), β -sitosterol (**5**), and chlorophyll a (**6**). The structures of **1-6** were identified by comparison of their ¹H and/or ¹³C NMR data with those reported in the literature.

INTRODUCTION

Artocarpus ovatus Blanco, an endemic Philippine tree, locally known as anubing is an evergreen tree growing up to 25 meters tall. A latex obtained from the tree has potential for use as a chewing gum base (Fern, 2012). The roasted seeds are eaten, while the timber which is strong and durable is used for construction (De la Cruz, 1991).

There is no reported study on the chemical constituents of *A. ovatus*. However, congeners of the tree have been studied for their chemical constituents and biological activities. There are about 50 species of the genus *Artocarpus* (Moraceae) which are sources of edible fruit and timber and are used in folk medicines. A review of the chemical constituents, biological and pharmacological activities of *Artocarpus* has been provided (Jagtap and Bapat, 2010). *Artocarpus* species are used for treatment against inflammation, malarial fever, diarrhoea, diabetes and tapeworm infection. They are rich in phenolic compounds such as flavonoids, stilbenoids, arylbenzofurans & jacalin (Jagtap and Bapat, 2010). Compounds from *Artocarpus* species exhibit diverse biological activities including antibacterial, antitubercular,

antiviral, antifungal, antiplatelet, antiarthritic, tyrosinase inhibitory and cytotoxicity (Jagtap and Bapat, 2010). This study is part of our research on the chemical constituents of the genus *Artocarpus* found in the Philippines.

We earlier reported the isolation of friedelinol, squalene, β -sitosterol, stigmasterol and phytol from the leaves of *A. camansi*, while the stems yielded polyprenol, cycloartenol and cycloartenol acetate (Tsai *et al.*, 2013). In another study, the leaves of *A. altilis* yielded β -sitosterol, triglycerides, squalene, polyprenol, lutein and fatty acids, while *A. odoratissimus* afforded β -sitosterol, triglycerides and fatty acids from the flesh of the fruit and seeds; and β -sitosterol, fatty acids and hydrocarbons from the fruit rind (Ragasa *et al.*, 2014). Furthermore, the unripe fruit of *A. heterophyllus* afforded cycloartenone, cycloartenol, and a diastereomeric mixture of 2,3-butanediols (Ragasa and Jorvina, 2004).

We report herein the fractionation by silica gel chromatography of the dichloromethane extract of the twigs of *A. ovatus*. The following compounds from the twigs were identified by NMR spectroscopy: lupeol (**1a**), α -amyrin (**1b**), β -amyrin (**1c**), lupeol fatty acid ester (**2a**), α -amyrin fatty acid ester (**2b**), β -amyrin fatty acid ester (**2b**), betulin (**3a**), 3β , 28-dihydroxyolean-12-ene (**3b**), oleanolic acid (**4**), β -sitosterol (**5**), and chlorophyll a (**6**) (Fig. 1). To the best of our knowledge this is the first report on the identification of these compounds from *A. ovatus*.

* Corresponding Author

Consolacion Y. Ragasa, Chemistry Department, De La Salle University, 2401 Taft Avenue, Manila 1004 Philippines.

Email: consolacion.ragasa@dlsu.edu.ph




Fig. 1: Chemical constituents of the twigs of *Artocarpus ovatus*: lubeol (**1a**), α -amvrin (**1b**), β -amvrin (**1c**), lubeol fatty acid ester (**2a**), α -amvrin fatty acid ester

MATERIALS AND METHODS

General Experimental Procedure

NMR spectra were recorded on a Varian VNMRs spectrometer in CDCl_3 at 600 MHz for ^1H NMR and 150 MHz for ^{13}C NMR spectra. Column chromatography was performed with silica gel 60 (70-230 mesh). Thin layer chromatography was performed with plastic backed plates coated with silica gel F₂₅₄ and the plates were visualized by spraying with vanillin/ H_2SO_4 solution followed by warming.

Sample Collection

The sample was collected from Bataan, Philippines in October 2013. It was identified as *Artocarpus ovatus* Blanco at the Jose Vera Santos Herbarium, Institute of Biology, University of the Philippines, Diliman, Quezon City.

General Isolation Procedure

A glass column 18 inches in height and 1.0 inch internal diameter was packed with silica gel. The crude extract from the twigs were fractionated by silica gel chromatography using increasing proportions of acetone in dichloromethane (10% increment) as eluents. Fifty milliliter fractions were collected.

All fractions were monitored by thin layer chromatography. Fractions with spots of the same *R_f* values were combined and rechromatographed in appropriate solvent systems until TLC pure isolates were obtained. A glass column 12 inches in height and 0.5 inch internal diameter was used for the rechromatography. Two milliliter fractions were collected. Final purifications were conducted using Pasteur pipettes as columns. One milliliter fractions were collected.

Isolation

The air-dried twigs of *A. ovatus* (278.7 g) were ground in a blender, soaked in CH_2Cl_2 for 3 days and then filtered. The solvent was evaporated under vacuum to afford a crude extract (8.1 g) which was chromatographed using increasing proportions of acetone in CH_2Cl_2 at 10% increment. The 10% acetone in CH_2Cl_2 fraction was rechromatographed (5 \times) in 5% EtOAc in petroleum ether to afford a mixture of **2a-2c** (5 mg) after washing with petroleum ether. The 20% acetone in CH_2Cl_2 fraction was rechromatographed (4 \times) in 15% EtOAc in petroleum ether to afford a mixture of **1a-1c** (1 mg) after washing with petroleum ether. The 30% acetone in CH_2Cl_2 fraction was rechromatographed

(3 ×) in 20% EtOAc in petroleum ether to afford **5** (6 mg) after washing with petroleum ether. The 40% acetone in CH₂Cl₂ fraction was rechromatographed (3 ×) in 20% EtOAc in petroleum ether to afford a mixture of **3a** and **3b** (2 mg) after washing with petroleum ether. The 50% acetone in CH₂Cl₂ fraction was rechromatographed (4 ×) in CH₃CN:Et₂O:CH₂Cl₂ (1:1:8) by volume ratio to afford **4** (4 mg) after washing with petroleum ether. The 60% acetone in CH₂Cl₂ fraction was rechromatographed (4×) in CH₃CN:Et₂O:CH₂Cl₂ (1:1:8) by volume ratio to afford **6** (5 mg) after washing with petroleum ether, followed by Et₂O.

Lupeol (1a)

colorless solid. ¹H NMR (600 MHz, CDCl₃): ¹H NMR (CDCl₃, 500 MHz) δ 4.68 (H-29b), 4.55 (H-29a), 3.18 (H-3), 1.68 (s, H₃-30), 0.96 (s, H₃-23), 0.78 (s, H₃-24), 0.83 (s, H₃-25), 0.94 (s, H₃-26), 1.06 (s, H₃-27), 0.91 (s, H₃-28), 1.68 (s, H₃-30).

α-Amyrin (1b)

Colorless solid, ¹H NMR (600 MHz, CDCl₃): ¹H NMR (CDCl₃, 500 MHz) δ 3.15 (H-3), 0.67 (H-5), 5.15 (H-12), 0.94 (s, H₃-23), 0.76 (s, H₃-24), 0.75 (s, H₃-25), 0.89 (s, H₃-26), 1.02 (s, H₃-27), 0.94 (s, H₃-28), 0.85 (d, *J* = 6.0 Hz, H₃-29), 0.74 (d, *J* = 7.0 Hz, H₃-30).

β-Amyrin (1c)

Colorless solid. ¹H NMR (600 MHz, CDCl₃): ¹H NMR (CDCl₃, 500 MHz) δ 3.15 (H-3), 0.67 (H-5), 5.22 (H-12), 0.77 (s, H₃-23), 0.90 (s, H₃-24), 0.74 (s, H₃-25), 0.94 (s, H₃-26), 1.16 (s, H₃-27), 1.06 (s, H₃-28), 0.86 (s, H₃-29), 0.79 (s, H₃-30).

Lupeol fatty acid ester (2a)

Colorless solid. ¹³C NMR (150 MHz, CDCl₃): 38.62 (C-1), 23.42 (C-2), 80.49 (C-3), 38.61 (C-4), 55.48 (C-5), 18.23 (C-6), 34.71 (C-7), 40.97 (C-8), 50.99 (C-9), 38.62 (C-10), 21.06 (C-11), 25.17 (C-12), 38.62 (C-13), 42.64 (C-14), 27.49 (C-15), 35.49 (C-16), 43.95 (C-17), 48.36 (C-18), 47.90 (C-19), 150.89 (C-20), 29.36 (C-21), 39.94 (C-22), 27.49 (C-23), 16.93 (C-24), 14.82 (C-27), 18.23 (C-28), 109.39 (C-29), 19.35 (C-30), 173.69 (C-1'), 34.87 (C-2'), 31.92 (C-3'), 22.69, 25.17, 29.18-29.70 (CH₂)_n, 14.12 (CH₃).

α-Amyrin fatty acid ester (2b)

Colorless solid. ¹³C NMR (150 MHz, CDCl₃): 38.68 (C-1), 22.69 (C-2), 80.49 (C-3), 37.14 (C-4), 55.48 (C-5), 18.37 (C-6), 32.48 (C-7), 39.68 (C-8), 47.89 (C-9), 37.11 (C-10), 23.31 (C-11), 124.66 (C-12), 138.73 (C-13), 42.17 (C-14), 28.70 (C-15), 26.56 (C-16), 33.75 (C-17), 59.04 (C-18), 39.59 (C-19), 39.68 (C-20), 31.25 (C-21), 41.51 (C-22), 28.06 (C-23), 16.92 (C-26), 23.42 (C-27), 28.06 (C-28), 17.46 (C-29), 21.38 (C-30), 173.69 (C-1'), 34.87 (C-2'), 31.92 (C-3'), 22.69, 25.17, 29.18-29.70 (CH₂)_n, 14.12 (CH₃).

β-Amyrin fatty acid ester (2c)

Colorless solid. ¹³C NMR (CDCl₃): 38.62 (C-1), 22.69

(C-2), 80.49 (C-3), 37.14 (C-4), 55.48 (C-5), 18.37 (C-6), 32.48 (C-7), 39.68 (C-8), 47.22 (C-9), 37.14 (C-10), 23.66 (C-11), 121.95 (C-12), 144.30 (C-13), 41.51 (C-14), 26.08 (C-15), 26.91 (C-16), 32.48 (C-17), 47.83 (C-18), 46.76 (C-19), 31.25 (C-20), 34.87 (C-21), 37.14 (C-22), 28.06 (C-23), 16.92 (C-24), 16.92 (C-26), 26.08 (C-27), 28.06 (C-28), 33.31 (C-29), 23.66 (C-30), 173.69 (C-1'), 34.87 (C-2'), 31.92 (C-3'), 22.69, 25.17, 29.18-29.70 (CH₂)_n, 14.12 (CH₃).

Betulin (3a)

Colorless solid. ¹H NMR (600 MHz, CDCl₃): δ ¹H NMR (CDCl₃, 500 MHz) δ 3.18 (H-3), 0.96 (s, H₃-23), 0.74 (s, H₃-24), 0.81 (s, H₃-25), 0.98 (s, H₃-26), 1.00 (s, H₃-27), 3.33 (d, H-28a), 3.79 (d, H-28b), 4.55 (d, H-29a), (d, 4.68 H-29b), 1.66 (s, H₃-30).

3β, 28-Dihydroxyolean-12-ene (3b)

Colorless solid. ¹H NMR (600 MHz, CDCl₃): δ ¹H NMR (CDCl₃, 500 MHz) δ 3.18 (H-3), 5.18 (d, H-12), 0.96 (s, H₃-23), 0.74 (s, H₃-24), 0.87 (s, H₃-25), 0.98 (s, H₃-26), 1.15 (s, H₃-27), 0.95 (H₃-29), 0.88 (H₃-29), 3.20 (d, H-28a), 3.52 (d, H-28b).

Oleanolic acid (4)

Colorless solid. ¹H NMR (600 MHz, CDCl₃): δ ¹H NMR (CDCl₃, 500 MHz) δ 3.20 (dd, *J* = 4.2, 11.4 Hz, H-3α), 5.26 (t, *J* = 3.6 Hz, H-12), 2.81 (dd, *J* = 4.2, 13.8 Hz, H-18), 0.96 (s, H₃-23), 0.73 (s, H₃-24), 0.89 (s, H₃-25), 0.75 (s, H₃-26), 1.11 (s, H₃-27), 0.91 (s, H₃-29), 0.88 (s, H₃-30); ¹³C NMR (150 MHz, CDCl₃): δ 38.38 (C-1), 27.17 (C-2), 77.02 (C-3), 38.74 (C-4), 55.20 (C-5), 18.28 (C-6), 32.59 (C-7), 39.25 (C-8), 47.62 (C-9), 37.07 (C-10), 23.38 (C-11), 122.63 (C-12), 143.58 (C-13), 41.57 (C-14), 27.67 (C-15), 22.90 (C-16), 46.51 (C-17), 40.96 (C-18), 45.86 (C-19), 30.67 (C-20), 33.78 (C-21), 32.42 (C-22), 28.09 (C-23), 15.53 (C-24), 15.31 (C-25), 17.12 (C-26), 25.93 (C-27), 183.29 (C-28), 33.06 (C-29), 23.56 (C-30).

β-Sitosterol (5)

¹H NMR (600 MHz, CDCl₃): δ 3.51 (m, H-3), 2.26, 2.21 (H₂-4), 5.33 dd (*J* = 3.0, 2.4 Hz, H-6), 0.66 (s, H₃-18), 0.99 (s, H₃-19), 0.90 (d, *J* = 6.6 Hz, H₃-21), 0.79 (d, *J* = 7.2 Hz, H₃-26), 0.82 (d, *J* = 6.6 Hz, H₃-27), 0.85 (t, *J* = 7.8 Hz, H₃-29).

Chlorophyll a (6)

¹H NMR (600 MHz, CDCl₃): δ 3.39 (s, H-1), 7.98 (dd, *J* = 18, 12 Hz, H-2a), 6.26 (dd, *J* = 18, 1.2 Hz, H-2b), 6.26 (dd, *J* = 12, 1.2 Hz, H-2b), 3.22 (s, H-3a), 3.67 (s, H-4a), 1.69 (t, *J* = 7.2, H-4b), 3.76 (s, H-1α or H-5α), 4.17 (m, H-7), 2.15, 2.46 (m, H₂-7a), 2.32, 2.60 (m, H₂-7b), 4.42 (m, H-8), 1.78 (d, *J* = 7.2 Hz, H₃-8a), 6.24 (s, H-10), 3.86 (s, H₃-10b), 9.38 (s, H-α or H-β), 9.51 (s, H-α or H-β), 8.54 (s, H-δ), 4.46 (m, P1), 5.11 (t, *J* = 1.2 Hz, P2), 1.56 (br s, P17), 0.82 (d, *J* = 6.6 Hz, P18 and P19), 0.76 (d, *J* = 6.6 Hz, P16), 0.74 (d, *J* = 6.6 Hz, P20).

RESULTS AND DISCUSSION

Silica gel chromatography of the dichloromethane extract of the twigs of *A. ovatus* afforded lupeol (**1a**) (Prakash and Prakash), α -amyrin (**1b**) (Raga *et al.*, 2011a), β -amyrin (**1c**) (Raga *et al.*, 2011a), lupeol fatty acid ester (**2a**) (Raga *et al.*, 2011b), α -amyrin fatty acid ester (**2b**) (Miranda *et al.*, 2006), β -amyrin fatty acid ester (**2c**) (Barreiros *et al.*, 2012), betulin (**3a**) (Tijjani *et al.*, 2012), 3β , 28-Dihydroxyolean-12-ene (**3b**) (Rahman *et al.*, 2007), oleanolic acid (**4**) (Ragasa and Lim, 2005), β -sitosterol (**5**) (Cayme and Ragasa, 2004), and chlorophyll a (**6**) (Ragasa and de Jesus, 2014). The structures of **1-6** were identified by comparison of their ^1H and/or ^{13}C NMR data with those reported in the literature.

Although bioassays were not conducted on the isolated compounds, there were previous studies that reported on their biological activities. Lupeol (**1a**) exhibited antiurolithiatic and diuretic activity (Vidya *et al.*, 2002). It prevented the formation of vesical calculi and reduced the size of the preformed stones in rats (Anand *et al.* 1994). α -Amyrin (**1b**) and β -amyrin (**1c**) were reported to possess anti-inflammatory activity (Recio *et al.* 1995; Madeiros *et al.* 2007), while α -amyrin, β -amyrin, and the 3-O-acyl derivatives of α -amyrin (**2b**) and β -amyrin (**2c**) exhibited analgesic property (Otuki *et al.* 2005; Soldi *et al.* 2008).

Betulin (**3a**) inhibited cell growth and induction of apoptosis in human melanoma cell line (Orchel *et al.*, 2014). It also induces apoptosis of *Leishmania donovani*. Treatment of promastigotes with betulin led to mitochondrial membrane damage, activation of caspase-like proteases, and DNA fragmentation in *L. donovani* promastigotes (Saudagar and Dubey, 2014).

An earlier study reported that ursolic acid and oleanolic acid (**4**) exhibited anti-tumor activity against human colon carcinoma cell line HCT15 with ursolic acid showing stronger activity than oleanolic acid (Li *et al.*, 2002). It exhibited anti-inflammatory effects by inhibiting hyperpermeability, the expression of CAMs, and the adhesion and migration of leukocytes (Lee *et al.*, 2013). It showed anti-inflammatory activities through the inhibition of the HMGB1 signaling pathway (Yang *et al.*, 2012). It has anti-inflammatory, hepatoprotective, gastroprotective, anti-ulcer, and immunoregulatory effect (Vachalkova *et al.*, 2004), gastroprotective effect on experimentally induced gastric lesions in rats and mice (Astudillo *et al.*, 2002), inhibits mouse skin tumor (Oguro, 1998), protects against hepatotoxicants and is used in China to treat hepatitis (Lui *et al.*, 1993), and significant antitumor activity on human colon carcinoma cell line HCT 15 (Li *et al.*, 2002).

β -Sitosterol (**5**) was observed to have growth inhibitory effects on human breast MCF-7 and MDA-MB-231 adenocarcinoma cells (Awad *et al.*, 2007). It was shown to be effective for the treatment of benign prostatic hyperplasia (Jayaprakasha *et al.*, 2007). It was also reported to attenuate β -catenin and PCNA expression, as well as quench radical *in-vitro*, making it a potential anticancer drug for colon carcinogenesis (Baskar *et al.*, 2010). It can inhibit the expression of NPC1L1 in

the enterocytes to reduce intestinal cholesterol uptake (Jesch *et al.*, 2009). It was reported to induce apoptosis mediated by the activation of ERK and the downregulation of Akt in MCA-102 murine fibrosarcoma cells (Moon *et al.*, 2007).

Chlorophyll (**6**) and its various derivatives are used in traditional medicine and for therapeutic purposes (Edwards, 1954). Natural chlorophyll and its derivatives have been studied for wound healing (Kephart, 1955), anti-inflammatory properties (Larato and Pfao, 1970), control of calcium oxalate crystals (Tawashi *et al.*, 1980), utilization as effective agents in photodynamic cancer therapy (Sternberg *et al.*, 1998; Nourse *et al.*, 1988; Henderson *et al.*, 1997), and chemopreventive effects in humans (Egner *et al.*, 2001; Egner *et al.*, 2003). A review on digestion, absorption and cancer preventive activity of dietary chlorophyll has been provided (Hardwick *et al.*, 1997).

Thus, the dichloromethane extract of the twigs of *Artocarpus ovatus* afforded compounds (**1-6**) with diverse biological activities.

REFERENCES

- Anand R, Patnaik GK, Kulshreshtha DK, Dhawan N. Antiurolithiatic activity of lupeol, the active constituent of *Crateva nuriala*. *Phytother Res* 1994; 8(7):417-421.
- Awad AB, Chinnman M, Fink CS, Bradford PG. β -Sitosterol activates Fas signaling in human breast cancer cells. *Phytomed* 2007; 14:747-754.
- Barreiros ML, David JM, Pereira PA, Guedes MLS, David JP. (2002). Fatty acid esters of triterpenes from *Erythroxylum passerinum*. *J Braz Chem Soc* 2002; 13:669-673.
- Baskar AA, Ignacimuthu S, Paulraj G, Numair K. Chemopreventive potential of β -sitosterol in experimental colon cancer model - an *In vitro* and *In vivo* study. *BMC Comp Alt Med* 2010; 10: 24.
- Cayme J-MC, Ragasa CY. Structure elucidation of stigmaterol and β -sitosterol from *Sesbania grandiflora* Linn. Pers. and β -Carotene from *Heliotropium indicum* Linn. by NMR Spectroscopy. *Kimika* 2004; 20(1/2):5-12.
- De la Cruz Jr FS. *Artocarpus odoratissimus* Blanco. 1991. *In Plant Resources of South-East Asia. 2. Edible Fruits and Nuts*. E.W.M. Verheij and R.E. Coronel (eds.). PROSEA, Pudoc, Wageningen. 94-96.
- Edwards BJ. Treatment of chronic leg ulcers with ointment containing soluble chlorophyll. *Physiother* 1954; 40:177-179.
- Egner PA, Munoz A, Kensler TW. Chemoprevention with chlorophyllin in individuals exposed to dietary aflatoxin. *Mutat Res* 2003; 52(3):209-216.
- Egner PA, Wang JB, Zhu YR, Zhang BC, Wu Y, Zhang QN. Chlorophyllin intervention reduces aflatoxin-DNA adducts in individuals at high risk for liver cancer. *Proc Natl Acad Sci* 2001; 98(25):1401-1406.
- Fern K. 2012. Useful Tropical Plants Database.
- Hardwick SJ, Carpenter KH, Law NS, Van Der Veen C, Marchant CE, Hird R, Mitchinson MJ. *Free Radic Res* 1997; 26 (4):351-362.
- Henderson BW, Bellnier DA, Greco WR, Sharma A, Pandry RK, Vaughan LA. An *in Vivo* Quantitative Structure-Activity Relationship for a Congeneric Series of Pyropheophorbide Derivatives as Photosensitizers for Photodynamic Therapy. *Cancer Res* 1997; 57:4000-4007.
- Jagtap UB, Bapat VA, Artocarpus: a review of its traditional uses, phytochemistry and pharmacology. *J Ethnopharmacol* 2010; 129(2):142-66.
- Jayaprakasha GK, Mandadi KK, Poulouse SM, Jadegoud Y, Gowda GA, Patil BS. Inhibition of colon cancer growth and antioxidant activity of bioactive compounds from *Poncirus trifoliata* (L.) Raf. *Bioorg Med Chem*, 2007; 15:4923-4932.

Jesch ED, Seo JM, Carr TP, Lee JY. Sitosterol reduces messenger RNA and protein expression levels of Niemann-Pick C1-like 1 in FHs 74 Int cells. *Nutr Res* 2009; 29(12):859-66.

Kephart JC. Chlorophyll derivatives - their chemistry, commercial preparation and uses. *Econ Bot* 1955; 9:3-18.

Larato DC, Pfao FR. Effects of a water-soluble chlorophyllin ointment on gingival inflammation. *N Y State Dent J* 1970; 36:291-293.

Lee W, Yang EJ, Ku SK, Song KS, Bae JS. Anti-inflammatory effects of oleanolic acid on LPS-induced inflammation *in-vitro* and *in-vivo*. *Inflammation* 2013; 36(1):94-102.

Li J, Guo W-J, Yang Q-Y. Effects of ursolic acid and oleanolic acid on human colon carcinoma cell line HCT15. *World J Gastroenterol* 2002; 8(3):493-495.

Madeiras R, Otuki MF, Avellar MC, Calixto JB. 2007. Mechanisms underlying the inhibitory actions of the pentacyclic triterpene -amyrin in the mouse skin inflammation induced by phorbol ester 12-O-tetradecanoylphorbol-13-acetate. *Eur J Pharmacol* 55(9):227-235.

Miranda RRS, Silva GDF, Duarte LP, Fortes ICP, Vieira Filho SA. Structural determination of 3 β -stearoyloxy-urs-12-ene from *Maytenus salicifolia* by 1D and 2D NMR and quantitative ¹³C NMR spectroscopy. *Magnetic Resonance in Chemistry* 2006; 44:127-131.

Moon DO, Kyeong Jun L, Yung HC, Gi-Young K. Moon DO, Kyeong Jun L, Yung HC, Gi-Young K. *Int Immunopharmacol* 2007; 7:1044-1053.

Nourse WL, Parkhurst RM, Skinner WA, Jordan RT. Photodynamic toxicity of porphyrins and chlorins for a human tumor cell line: combined light and concentration dose responses for the retained fraction. *Biochem Biophys Res Commun* 1988; 151:506-511.

Orchel A, Kulczycka A, Chodurek E, Bebenek E, Borkowska P, Boryczka S, Kowalski J, Dzierzewicz X. Influence of betulin and 28-O-propynylbetulin on proliferation and apoptosis of human melanoma cells (G-361). *Postepy Hig Med Dosw*, 2014; 68(0):191-197.

Otuki C, Ferreira J, Lima F, Meyre-Silva C, Malheiros A, Muller L, Cani G, Santos A, Yunes R, Calixto J. Antinociceptive properties of a mixture of α -amyrin and β -amyrin triterpenes: evidence for participation of protein kinase C and protein kinase A pathways. *J Pharmacol Exp Therapeutics* 2005; 31(1):310-318.

Prakash CVS, Prakash I. Isolation and Structural Characterization of Lupane Triterpenes from *Polypodium vulgare*. *Res J Pharm Sci* 2012; 1(1):23-27.

Raga DD, Alimboyoguen AB, Shen C-C, Ragasa CY. Triterpenoids and an anti-angiogenic sterol from *Ardisia pyramidalis* Cav. *Pers. Philipp Agric Sci* 2011a; 94(2):103-110.

Raga DD, Cheng CLC, Lee KCIC, Olaziman WZP, De Guzman VJA, Shen C-C, Franco FC, Ragasa CY. Bioactivities of triterpenes and sterol from *Syzygium samarangense*. *Z Naturforsch* 2011b; 66c:235-244.

Ragasa CY, Ng VA, Park JH, Kim DW, Cornelio K, Shen C-C. Chemical Constituents of *Artocarpus altilis* and *Artocarpus odoratissimus*. *Res J Pharm Biol Chem Sci* 2014; 5(4):1081-1087.

Ragasa CY, Jorvina K, Rideout JA. Antimicrobial compounds from *Artocarpus heterophyllus*. *Philipp J Sci* 2004; 133(2):97-101.

Ragasa CY, Lim K. Secondary metabolites from *Schefflera odorata* Blanco. *Philipp J Sci* 2005; 134(1):63-67.

Ragasa CY, de Jesus J. Porphyrins and polyprenols from *Macaranga tanarius*. *Res J Pharm Biol Chem Sci* 2014; 5(3):701-708.

Rahman MZ, Sultana SJ, Faruquee CF, Ferdous F, Rahman MS, Islam MS, Rashid MA. Phytochemical and biological investigations of *Erythrina variegata*. *Saudi Pharm J* 2007; 15(2):140-145.

Recio MC, Giner RM, Manez S, Rios JL. Structural requirements for the anti-inflammatory activity of natural triterpenoids. *Planta Med* 1995; 61(2):181-185.

Saudagar P, Dubey VK. Molecular mechanisms of *in vitro* betulin-induced apoptosis of *Leishmania donovani* Am J Trop Med Hyg 2014; 90(2):354-360.

Soldi C, Pizzolatti G, Luiz A, Marcon R, Meotti F, Miotob L, Santos A. Synthetic derivatives of the α - and β -amyrin triterpenes and their antinociceptive properties. *Bioorganic and Medicinal Chemistry* 2008; 16(6):3377-3386.

Sternberg ED, Dolphin D, Bruckner C. Porphyrin-based Photosensitizers for Use in Photodynamic Therapy. *Tetrahedron* 1998; 54:4151-4152.

Tawashi R, Cousineau M, Sharkawi M. Effect of sodium copper chlorophyllin on the formation of calcium oxalate crystals in rat kidney. *Invest Urol* 1980; 18:90-92.

Tijjani A, Ndukwe IG, Ayo RG. Isolation and characterization of lup-20(29)-ene-3,28-diol (betulin) from the stem-bark of *Adenium obesum* (Apocynaceae). *Trop J Pharm Res* 2012; 11 (2):259-262.

Tsai P-W, de Castro-Cruz K, Shen C-C, Ragasa CY. Chemical constituents of *Artocarpus camansi*. *Phcog J* 2013; 5:80-82.

Vidya L, Leni M, Varalakshmi P. 2002. Evaluation of the effect of triterpenes on urinary risk factors of stone formation in pyridoxine hyperoxaluric rats. *Phytother Res* 16 (6):514-518.

Yang EJ, Lee W, Ku SK, Song KS, Bae JS. Anti-inflammatory activities of oleanolic acid on HMGB1 activated HUVECs. *Food Chem Toxicol* 2012; 50(5) 1288-94.

How to cite this article:

Consolacion Y. Ragasa, Jocelyn L. Caro, Chien-Chang Shen. Triterpenes and Sterol from *Artocarpus ovatus*. *J App Pharm Sci*, 2014; 4 (10): 007-011.