



Received: 23-04-2011
 Revised on: 26-04-2011
 Accepted: 03-05-2011

Comparison effect of Pioglitazone and Glimepiride alone on renal function marker in experimentally induced renal damage in diabetic rats

Jagdish Kakadiya and Nehal Shah

ABSTRACT

Present study was designed to evaluate in Renoprotective activity of Pioglitazone and Glimepiride on Ischemia/reperfusion (I/R) induced renal damage in diabetic rats. Ischemia/reperfusion injury, which is commonly seen in the field of renal surgery or transplantation in diabetic condition, is a major cause of acute renal failure. Type 2 Diabetes was induced in rats by a single intraperitoneal (i.p) injection of Streptozotocin (65 mg/kg, STZ) in overnight fasting rats followed by the i.p administration of Nicotinamide (110 mg/kg, NIC) after 15 minutes. After right nephrectomy, Pioglitazone (10 mg/kg/day, p.o) and Glimepiride (0.5 mg/kg/day, p.o) were administered for 15 days. On the 16th day, ischemia was induced in contra lateral kidney for 45 min, followed by reperfusion for 24 hr. renal function marker and oxidative parameter were estimated at the end of 24 hr reperfusion. At the end of experimental period the level of malondialdehyde formation/ lipid peroxidation (LPO) in kidney tissue and serum marker Creatinine, Urea and Uric acids were significantly increased. Whereas, the activity of biomarkers of oxidative stress such as reduced glutathione (GSH), Catalase (CAT) and superoxide dismutase (SOD) were found to be decreased significantly compared to control rats. Pioglitazone improved the renal dysfunction and oxidative stress after renal ischemia/reperfusion injury in diabetic rats, but Glimepiride less improved the renal marker change compared to treatment of Pioglitazone. In conclusion, Pioglitazone shows potent may improve renal function marker and oxidative stress in kidney in I/R induced renal damage in type 2 diabetic rats.

Key words: Pioglitazone; renal ischemia reperfusion injury; Renal marker; type 2 diabetes

INTRODUCTION

Ischemic injury to brain, heart, and kidney is associated with high morbidity and mortality. Improving the ability of these organs to tolerate ischemic injury would have important implications. Ischemic insults are often recurrent in diabetic patients. In the setting of loss of renal blood flow autoregulation that characterizes the post-ischemic kidney (Conger, 1997), Renal ischemia/reperfusion (I/R) injury is a major cause of acute renal failure (ARF) (Radhakrishnan et al., 1997), which is faced in many clinical situations such as kidney transplantation, partial nephrectomy, renal artery, angioplasty, aortic aneurysm surgery, and elective urological operations. In these conditions, I/R injury initiates a complex and interrelated sequence of events, resulting in injury to and the eventual death of renal cells (Thadhani et al., 1996; Paller, 1998). Several factors have been implicated in the pathophysiological changes occurring while renal I/R injury including vascular or microvascular injury, endothelial dysfunction, accelerated cell necrosis, granulocyte activation, and modulation of nitric oxide/angiotensin II axis (Maxwell et al., 1997; Adam et al., 2000).

The rennin-angiotensin system plays a pivotal role in regulation of blood pressure. Renin acts on angiotensinogen to form angiotensin I, which is converted to angiotensin II with the help of angiotensin-converting enzyme (ACE) (Gavras et al., 1997). Angiotensin II is an important mediator in kidney injury. Accumulating evidence suggests that angiotensin II stimulates intracellular formation of reactive oxygen species (ROS) such as the superoxide anion and

Jagdish Kakadiya

Pharmacy Department, Jodhpur
 National University, Narnadi,
 Jhanwar road, Jodhpur, Rajasthan,
 India.

Nehal Shah

Pharmacy Department, Dharmaj Degree
 Pharmacy College, Petlad -Khambhat
 Road, Dharmaj, Anand, India.

*For Correspondence:

Mr. Jagdish L. Kakadiya
 Pharmacology department,
 Dharmaj Degree Pharmacy College,
 Dharmaj, Anand, Gujarat, India.
 Email: jagdishkakadiya@yahoo.co.in

(Gumieniczek, 2003) has been reported. Pioglitazone (PIO) hydrochloride is a widely used drug in the treatment of insulin resistance diabetes. PIO showed dose dependant beneficial effects in many of the pathological conditions including reduction in blood glucose, lowering blood pressure and restoring endothelial functions in animals (Jayesh et al., 2005). Pioglitazone lowers blood pressure and restores blunted endothelium-dependent vasodilatation in fructose-fed rats (Kotchen et al., 1997), insulin-resistant rhesus monkey (Kemnitz et al., 1994), SHR (Grinsell et al., 2000) and sucrosefed SHR (Uchida et al., 1997). Glimperide (GLI) an oral blood glucose lowering drug of the sulfonylurea class is reported to have pancreatic and extra pancreatic effects as well. The blockages of K_{ATP} channels of pancreatic cells by sulphonylurea are critical in the regulation of glucose regulated insulin secretion.

It has been demonstrated to be effective in the treatment of non-insulin dependent diabetes mellitus with insulin resistance. GLI is pharmacologically distinct from glibenclamide because of differences in receptor-binding properties which could result in a reduced binding to cardiomyocyte K_{ATP} channels. Some have reported the antioxidative properties of GLI. Moreover, GLI is pharmacologically different than other sulfonylurea drugs for its cardiovascular effects. The present study, we investigated the protective effect of Pioglitazone and Glimperide on renal I/R injury in diabetic rats and other word effect of Pioglitazone alone and Glimperide on renal marker and oxidative stress of kidney tissues in I/R induced renal damage in diabetic rats.

MATERIALS AND METHODS

Drugs and Chemicals

Pioglitazone hydrochloride and Glimperide was obtained as a gift sample from Alembic Pharmaceuticals Pvt. Ltd., Baroda, India. STZ and NIC were obtained from SIGMA, St. Louis, MO, USA. All other chemicals and reagents used in the study were of analytical grade.

Experimental Animals

All experiments and protocols described in present study were approved by the Institutional Animal Ethics Committee (IAEC) of Dharmaj Degree Pharmacy College, Anand. Sprague Dawley rats (210 ± 15 g) were housed in-group of 3 animals per cage and maintained under standardized laboratory conditions (12-h light/dark cycle, 24°C) and provided free access to palletted CHAKKAN diet (Nav Maharashtra Oil Mills Pvt., Pune) and purified drinking water *ad libitum*. The animal experiment was approved by Animal Ethical Committee of the Institute (1163/a/08/CPCSEA).

Experimental Induction of Type 2 Diabetes in Rats

Type 2 Diabetes was induced in rats by a single intraperitoneal (i.p) injection of Streptozotocin (65 mg/kg, STZ) in overnight fasting rats or mice followed by the i.p administration of Nicotinamide (110 mg/kg, NIC) after 15 minutes. STZ was dissolved in citrate buffer (pH 4.5) and NIC was dissolved in

normal saline. After 7 days following STZ and NIC administration, blood was collected from retro-orbital puncture and serum samples were analyzed for blood glucose (Masiello et al., 1998). Animals showing fasting blood glucose higher than 250 mg/dL were considered as diabetic and used for the further study.

Experimental Protocol

The rats were divided into three groups each consisting of six animals:

Group 1: Animals served as sham-operated (underwent all surgical procedures without ischemia reperfusion).

Group 2: After right nephrectomy on day 1, vehicle (0.5 % sodium CMC) was administered for 15 days; on day 16, ischemia was produced in the left kidney for 45 min, followed by reperfusion of 24 hr (I/R control).

Group 3: After right nephrectomy on day 1, Pioglitazone (10 mg/kg/day, p.o.) was administered for 15 days; on day 16, ischemia was produced in the left kidney for 45 min, followed by reperfusion of 24 hr (I/R + PIO).

Group 4: After right nephrectomy on day 1, Glimperide (0.5 mg/kg/day, p.o.) was administered for 15 days; on day 16, ischemia was produced in the left kidney for 45 min, followed by reperfusion of 24 hr (I/R + GLI).

Surgical Procedure

The progress of the experiment		
	Unilateral	right
Day 1	nephrectomy	
Day 15	Treatment	
Day 16	45 minutes ischemia (left kidney)	
Day 17	24 hr reperfusion	

Right nephrectomy was performed through a right flank incision (2 cm) under general anesthesia, ketamine (100 mg/kg, i.p.). After right nephrectomy, several treatments were given as mentioned previously for 15 days. On day 16, ischemia was produced in the left kidney by performing a left flank incision and dissecting the left renal pedicle to expose the renal vessels. Non traumatic vascular clamps were used to stop blood flow (in artery and vein) for 45 min. Reperfusion was established by removing the clamp after 45 min ischemia. The abdominal wall (muscular layer and skin) was closed with 4.0 mononylon suture. At the end of reperfusion period (after 24 hr), blood samples were collected and used for the estimation of renal function (BUN and creatinine). The abdomen was opened, and the kidneys were harvested for the biomarkers of oxidative stress.

Characterization of Type 2 Diabetes Model

Type 2 diabetes was confirmed by measuring fasting serum glucose using standard diagnostic kit (SPAN diagnostics Pvt., India) and the degree of uncontrolled diabetic state was confirmed by measuring HbA1c (Ion Exchange Resin method). After seven day, diabetes was confirmed by measuring glucose and HbA1c as mentioned above.

Estimation of kidney function marker

Blood was collected from the rats by retro-orbital puncture at the time of sacrifice and was allowed to clot for 10 minutes at room temperature. Clots were centrifuged at 2500 rpm for 10 minutes to separate the serum. Serum creatinine and urea levels were measured by assay kits (SPAN Diagnostics Pvt. India) and Serum Uric acid levels were measured by assay kits (Crest Biosystems Ltd. India).

Preparation of Tissue Homogenate

After sacrificing the animals, their kidneys were quickly removed, perfused immediately with ice cold hypertonic saline solution, weighed and homogenized in chilled Tris buffer (10 mM, pH 7.4) at a concentration of 10% (w/v). The homogenates were centrifuged at 10,000×g at 0°C for 20 min using Remi C-24 high speed cooling centrifuge. The clear supernatant was used for the assay of following antioxidant parameters. The levels of Lipid peroxidation (LPO) formation and the activities of endogenous antioxidant enzymes such as catalase (CAT), reduced glutathione (GSH) and superoxide dismutase (SOD) were estimated by the method of Slater and Sawyer (Slater et al., 1971) Hugo Aebi as given by Hugo (Hugo et al., 1984) Moron et al (Moron et al., 1979) and Mishra and Fridovich (Mishra et al., 1992).

Statistical Analysis

All of the data are expressed as mean ± SEM. Statistical significance between more than two groups was tested using one-way ANOVA followed by the Bonferroni multiple comparisons test or unpaired two-tailed student's t-test as appropriate using a computer-based fitting program (Prism, Graphpad 5). Differences were considered to be statistically significant when $p < 0.05$.

RESULTS

Characterization of Type 2 Diabetes

Single intraperitoneal (i.p) injection of Streptozotocin (65mg/kg) followed by i.p administration of Nicotinamide (110 mg/kg) to rats produced severe hyperglycemia and increased HbA1c in 70 to 80 % the animals (Table 1).

Table 1. Effect of Streptozotocin (65mg/kg/day, p.o) and Nicotinamide (110 mg/kg/day, p.o) on serum glucose and HbA1c changes level in rats.

Groups	Glucose	HbA1c
CON	101.8 ± 6.799	5.455 ± 0.3729
STZ + NIC	332.8 ± 9.167***	9.900 ± 0.6323***

Values are expressed as mean ± SEM for six animals in the group. *** $P < 0.001$ considered statistically significant as compared to respective Control group.

Effect of PIO and GLI on kidney function marker

The six rats which underwent renal I/R exhibited a significant increase in the serum concentrations of creatinine ($P < 0.001$), urea ($P < 0.001$), and uric acid ($P < 0.001$) compared with the sham control animals, suggesting a significant degree of glomerular dysfunction mediated by renal I/R. In I/R+PIO treated diabetic rats, serum creatinine, urea and uric acid levels were significantly ($p < 0.05$, $n = 6$) higher as compared to I/R control group alone (Fig.1).




Figure 1. Effect of Pioglitazone (10 mg/kg/day, p.o) on serum Creatinine (A), Urea (B), and Uric acid (C) in the diabetic rats exposed to renal ischemia/reperfusion (I/R) injury. Values are expressed as mean ± SEM for six animals in the group. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ considered statistically significant as compared to respective Sham group.

Treatment with Glimperide significantly ($P < 0.05$) reduced Serum uric acid in I/R + GLI group compared to Sham group (Fig. 2). but, In I/R+GLI treated diabetic rats, serum creatinine and urea levels were significantly no change as compared to I/R control group alone (Fig.2).




Figure 2 Effect of Glimperide (0.5 mg/kg/day, p.o) on serum Creatinine (A), Urea (B), and Uric acid (C) in the diabetic rats exposed to renal ischemia/reperfusion (I/R) injury. Values are expressed as mean ± SEM for ten animals in the group. ns = Non Significant, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ considered statistically significant as compared to respective Sham group.

Effect of PIO and GLI on antioxidant activity

Renal I/R group of diabetic rats showed significantly decreased enzymatic activity of superoxide dismutase ($P < 0.001$), Catalase ($P < 0.001$), and reduced glutathione ($P < 0.001$) when compared with the sham control rats. These declining trends were significantly ($P < 0.05$, $P < 0.01$, $P < 0.05$, respectively Sham control Group) decreased in the group treated with PIO compared with those in the I/R-only group (Fig. 3).




Figure 3. Effect of Pioglitazone (10 mg/kg/day, p.o) on Superoxide dismutase (A), Catalase(B), Reduced glutathione (C) and lipid peroxidation (D) in the diabetic rats exposed to renal ischemia/reperfusion (I/R) injury. Values are expressed as mean \pm SEM for six animals in the group. * P<0.05, **P<0.01, ***P<0.001 considered statistically significant as compared to respective Sham group.

Renal I/R produced a significant (P<0.001) increase in MDA levels in comparison with the sham operation in the rats. Treatment with PIO before renal I/R was associated with a significantly (P<0.05) lower MDA level than that in the rats that underwent only renal I/R (Fig. 4). Renal I/R produced a significant (P<0.001) increase in MDA levels in comparison with the sham operation in the rats. Treatment with Glimperide significantly (P<0.05) increased MDA in I/R + GLI group compared to Sham group (Fig. 4).




Figure 4 Effect of Glimperide (0.5 mg/kg/day, p.o) on Superoxide dismutase (A), Catalase(B), Reduced glutathione (C) and lipid peroxidation (D) in the diabetic rats exposed to renal ischemia/reperfusion (I/R) injury. Values are expressed as mean \pm SEM for six animals in the group. *P<0.05, ***P<0.001 considered statistically significant as compared to respective Sham group.

DISCUSSION

The present study was under taken with the objective of exploring evaluate Pioglitazone on renal marker in I/R induced renal damage in diabetic rats. The transient discontinuation of renal blood supply is encountered in many clinical situations such as kidney transplantation, partial nephrectomy, renal artery angioplasty, aortic aneurysm surgery, and elective urological

operations (Thadhani et al., 1996; Conger, Paller, 1998). This transient discontinuation causes renal I/R injury which results in decreased glomerular filtration and renal blood flow and increased urine output characterized by natriuresis and impaired concentrating ability. Much of this tubular and glomerular dysfunction has been postulated to occur during the reperfusion period following anoxia, and generation of ROS has been postulated as one of the major factors contributing to this reperfusion injury. In renal I/R injury, ROS are capable of reacting with lipids leading to lipid peroxidation of biological membranes, which in turn impacts enzymatic processes, such as ion pump activity, inhibiting transcription and repair of DNA. If lipid peroxidation remains unchecked, it will ultimately result in cell death (Chatterjee et al., 2000; Singh et al., 2004).

In our study, animals subjected to renal I/R demonstrated an increase in the renal MDA and at attenuated antioxidant enzymes pool. Lipid peroxidation and antioxidant enzymes are important indexes of oxidant injury (Muller et al., 2000). Demonstrations of lipid peroxidation as indexes for oxidative damage may help us better understand the effects of ROS on the cellular components (Ozyurt et al., 2001). Renal I/R-induced oxidative stress was associated with impaired kidney function, leading to a marked increase in serum creatinine, urea, and uric acid levels. Pretreatment with Pioglitazone prevented renal I/R-induced lipid peroxidation and protected the kidneys from severe attenuation of antioxidant enzymes activity in rats exposed to the renal I/R. Furthermore, the impaired kidney function was significantly improved by Pioglitazone. The mechanism of the protective effect of Pioglitazone on renal I/R injury can be explained by its antioxidant activity (Gumieniczek, 2003). Pretreatment with Glimperide prevented renal I/R-induced lipid peroxidation and protected the kidneys from severe attenuation of only one parameter antioxidant enzymes activity in rats exposed to the renal I/R. Furthermore, the low impaired kidney function was significantly improved by Glimperide. The rennin-angiotensin system plays a pivotal role in regulation of blood pressure. Renin acts on angiotensinogen to form angiotensin-I, which is converted to angiotensin-II with the help of angiotensin-converting enzyme. Accumulating evidence suggests that angiotensin-II stimulates intracellular formation of ROS such as superoxide anion and hydrogen peroxide that leads to kidney damage (Sachse, 2007). Generation of ROS has been postulated as one of the major factors contributing to this reperfusion injury. Oxidative stress can result from increased ROS production, and/or from decreased ROS scavenging capability. The ROS attach to the polyunsaturated fatty acids in the membrane lipids and result in peroxidation, which may lead to disorganization of cell structure and function. After reperfusion and reoxygenation, the imbalance between restoration of oxygen supply and mitochondrial respiratory function results in massive generation of superoxide anion in mitochondria (Muller et al., 2000; Ozyurt et al., 2001). Under these conditions, the defensive system, which is known as antioxidant or antioxidant enzymes, cannot prevent the escape of ROS, especially in mitochondria, and their effects on other intracellular sites. This cascade of events is known as reperfusion injury (Ozyurt et al.,

2001). In this study, renal I/R increased oxidative stress products including tissue MDA and depleted the antioxidant enzymes pool, as is evident from the declined activity of superoxide dismutase, catalase, and reduced glutathione. It can be speculated that pretreatment with Pioglitazone prevented renal I/R-induced lipid peroxidation and protected the kidneys from severe increasing of ROS products and depletion of superoxide dismutase and reduced glutathione in rats exposed to the renal I/R.

CONCLUSIONS

Our data support a role for Pioglitazone may better improve compare to Glimepiride because Pioglitazone reduced renal marker and oxidative stress in ischemia/reperfusion induced renal marker in diabetic rats.

DECLARATION OF INTEREST

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the manuscript.

REFERENCE

- Adam A, Raji L. Nitric oxide--angiotensin II axis in renal and cardiovascular injury. *J Nephrol.*, 2000; 13:211-20.
- Chatterjee PK, Cuzzocrea S, Brown PA, et al. Tempol, a membrane-permeable radical scavenger, reduces oxidant stress-mediated renal dysfunction and injury in the rat. *Kidney Int.*, 2000; 58:658-73.
- Conger, J. *Adv. Renal Replace. Ther.* 4, Suppl., 1997; 1: 25-37.
- Gavras HP, Salerno CM. The angiotensin II type 1 receptor blocker losartan in clinical practice: a review. *Clin Ther.*, 1996; 18:1058-67.
- Grinsell JW, Lardinois CK, Swislocki A, Gonzalez R, Sare JS, Michaels JR, et al. Pioglitazone attenuates basal and postprandial insulin concentrations and blood pressure in the spontaneously hypertensive rat. *Am J Hypertens*, 2000; 13:370-5.
- Gumieniczek A. Effect of the new thiazolidinedione-pioglitazone on the development of oxidative stress in liver and kidney of diabetic rabbits. *Life Sci*, 2003; 74:553-62.
- Hugo EB. Oxidoreductases acting on groups other than CHOH: catalase. In: Colowick SP, Kaplan NO, Packer L, editors. *Methods in Enzymology*, vol. 105. London 7 Academic Press, 1984; 121-5.
- Jayesh B. Majithiya, Arvind N. Paramar, R. Balaraman. Pioglitazone, a PPAR γ -agonist, restores endothelial function in aorta of Streptozotocin-induced diabetic rats; *Cardiovascular Research*, 2005; 66:150-161.
- Kemnitz JW, Elson DF, Roecker EB, Baum ST, Bergman RN, Meglasson MD. Pioglitazone increases insulin sensitivity, reduces blood glucose, insulin, and lipid levels, and lowers blood pressure in obese, insulin-resistant Rhesus monkeys. *Diabetes*, 1994; 43:204-11.
- Kotchen TA, Reddy S, Zhang HY. Increasing insulin sensitivity lowers blood pressure in the fructose-fed rat. *Am J Hypertens*, 1997; 10:1020-6.
- Masiello, P., Broca, C., Gross, R., Roye, M., Manteghetti, M., Hillaire-Buys, D., Novelli, M., Ribes, G. Experimental NIDDM: development of a new model in adult rats administered Streptozotocin and Nicotinamide. *Diabetes*, 1998; 47: 224-229.
- Maxwell SR, Lip GY. Reperfusion injury: a review of the pathophysiology, clinical manifestations and therapeutic options. *Int J Cardiol.*, 1997; 58:95-117.
- Mishra HP, Fridovich I. The role of superoxide anion in the autooxidation of epinephrine and a simple assay for superoxide dismutase. *J Biochem*, 1992; 247:3170-5.
- Moron MS, Depierre JW, Mannervik B. Levels of glutathione, glutathione reductase and glutathione S-transferase activities in rat lung and liver. *Biochim Biophys Acta*, 1997; 582:67-78.
- Muller DN, Dechend R, Mervaala EM, et al. NF-kappaB inhibition ameliorates angiotensin II-induced inflammatory damage in rats. *Hypertension*, 2000; 35:193-201.
- Ozyurt H, Irmak MK, Akyol O, Sogut S. Caffeic acid phenethyl ester changes the indices of oxidative stress in serum of rats with renal ischaemia-reperfusion injury. *Cell Biochem Funct.*, 2001; 19:259-63.
- Paller MS. Acute renal failure: controversies, clinical trials, and future directions. *Semin Nephrol.*, 1998; 18:482-9.
- Radhakrishnan J, Kiryluk K. Acute renal failure outcomes in children and adults. *Kidney Int.*, 1997; 69:17-9.
- Sachse A, Wolf G. Angiotensin II-induced reactive oxygen species and the kidney. *J Am Soc Nephrol.*, 2007; 18:2439-46.
- Singh D, Chander V, Chopra K. The effect of quercetin, a bioflavonoid on ischemia/reperfusion induced renal injury in rats. *Arch Med Res.*, 2004; 35:484-94.
- Slater TF, Sawyer BC. The stimulatory effects of carbon tetrachloride and other halogeno alkanes or peroxidative reactions in rat liver fractions in vitro. *Biochem J*, 1971; 123:805-14.
- Thadhani R, Pascual M, Bonventre JV. Acute renal failure. *N Engl J Med.*, 1996; 334:1448-60.
- Uchida A, Nakata T, Hatta T, Kiyama M, Kawa T, Morimoto S, et al. Reduction of insulin resistance attenuates the development of hypertension in sucrose-fed SHR. *Life Sci*, 1997; 61(4):455-64.