

Review on phytochemical constituents and pharmacological activities of genus *Galium*

Mayssaloune Ali Kansa^{1*} , Mohamed Ali Hijazi¹ , Abdalla El-Lakany^{1,2} , Maha Aboul-Ela^{1,2} 

¹Department of Pharmaceutical Sciences, Faculty of Pharmacy, Beirut Arab University, Beirut, Lebanon.

²Department of Pharmacognosy, Faculty of Pharmacy, University of Alexandria, Alexandria, Egypt.

ARTICLE HISTORY

Received on: 27/04/2024
Accepted on: 30/07/2024
Available Online: 05/09/2024

Key words:

Galium, Rubiaceae,
phytochemistry,
pharmacology,
biological activity.

ABSTRACT

Medicinal plants are a rich source of phytochemical constituents of diverse structures that are behind their various pharmacological effects. Plants of genus *Galium* named Bedstraws, family *Rubiaceae*, are distributed throughout North and South America, Europe, the northern U.S., southern Canada, and tropical Asia. More than 600 species have been used in traditional medicine for treating different ailments. Owing to the valuable published uses of these plant species, and the versatility of their bioactive metabolites such as *Galium verum*, *Galium aparine*, *Galium mollugo*, and *G. odoratum*, it was deemed interesting to summarize the previous studies done on this genus to discuss the pharmacological profile of all isolated classes of metabolites. The search was adopted using some essential keywords such as *Galium*, phytochemistry, pharmacology, and biological activity from journals and books in databases such as Scopus, Elton B. Stephens CO, ScienceDirect, Embase, security identifier, and Medline from 1995 to 2024. The results showed that species belonging to *Galium* have various pharmacological activities including antimicrobial, antioxidant, anti-cancer, immunomodulatory, and anti-inflammatory effects because they are rich in phenolic compounds, iridoid glycosides, anthraquinones, phytosterols, saponins, and essential oils. Accordingly, this review will stimulate the scientific community for further research and boost the discovery of novel bioactive compounds from various species belonging to this genus distributed worldwide.

INTRODUCTION

Family Rubiaceae is one of the biggest angiosperm families having a wide number of species distributed worldwide. Its presence in almost all biomes and its diversity make it one of the most valuable components of tropical vegetation [1]. It has been the focus of researchers' attention because of its great pharmacological use being rich in bioactive metabolites including triterpenes, indole alkaloids, iridoids, flavonoids, and anthraquinones [2]. Plants belonging to Family Rubiaceae have been widely used in folk medicine due to their antioxidant, antiviral, antibacterial, anti-inflammatory, and analgesic effects in addition to their activity on the

central nervous system and vascular diseases [3]. Moreover their economic importance mainly the *Coffea arabica* species belonging to the Ixoroideae subfamily rich in caffeine that acts as a vasoconstrictor, central nervous system stimulant, diuretic, bronchodilator, and anti-migraine [4]. As well as the quinine isolated from *Cinchona* species in the Cinchonoideae subfamily by Pelletier and Caventou was for 200 years the only antimalarial drug and is considered responsible for the discovery of synthetic medicines for treating this disease [5]. Other research has isolated bioactive compounds including anthraquinones, iridoid glycosides, flavonols, sterols, and triterpene from *Crucianella maritima* L. (Rubiaceae) growing in Egypt and highlighted the cytotoxic and antimicrobial activities of these isolates [6,7]. The taxonomic classification of the Rubiaceae family is complex and incomplete according to Robbrecht's classification [8]. The fruits are derived from the bilocular ovary having locules with many ovules. The fruits are of red, orange, blackish, or yellow color and have

*Corresponding Author
Mayssaloune Ali Kansa, Department of Pharmaceutical Sciences, Faculty of Pharmacy, Beirut Arab University, Beirut, Lebanon.
E-mail: mayssalounekansa@yahoo.com

ovoid, spherical, or ellipsoid forms, small in size mainly in Gardenieae belonging to the Ixoroideae subfamily that was subclassified to tribes based on fruit characteristics such as the distinction between berries and drupes [9]. Family Rubiaceae is considered one of the five richest flowering plants having 13,000 species divided into three subfamilies, 620 genera, and 43 tribes. They are found on all continents but are mainly in subtropical and tropical areas. However, the relationships between taxa are not yet resolved since there are indications of polyphyletic/para genera [10]. It is classified into four subfamilies: antirheoideae, rubioideae, ixoroideae, and cinchonoidae. Family Rubiaceae is cosmopolitan, and pantropical with only a few species distributed extra tropically, around half of the species are available in the Neotropics. They are adapted to every habitat even desertic and arid climates. The majority of species belonging to this family are in the Atlantic forests of Brazil, Andean cloud forests, and the Amazon Basin with the endemism centered mainly in the Greater Antilles and the Guyana Highlands in the Neotropics [1].

The genus *Galium* L. is the largest in the tribe Rubieae belonging to the subfamily Rubioideae of family (Rubiaceae Juss.), it has around 600 species, distributed in North and South America, extratropical Eurasia, Australia, Arctic highlands, subtropical to tropical climatic zones, and North America. In Ukraine, about 50 species of Bedstraws are found [11]. This genus is also named yogurt herb because it contains an enzyme giving it the traditional use to coagulate milk [12]. *Galium* L. plants (family Rubiaceae Juss.) are commonly used in folk medicine for treating different ailments mainly infections of the skin, genitourinary systems, and respiratory tract because they are rich in lipophilic complexes of potent antimicrobial activity, mainly against Gram-positive microorganisms and slightly less against Gram-negative bacteria and low toxicity [13]. They are also traditionally choleric, diuretic, and anti-inflammatory. The most commonly used remedy is the tincture “Tazalok TM” based on *Galium verum* and used for menstrual disorders. Based on the medicinal importance of *Galium* species, and since the previous studies on plants of genus *Galium* have not completely covered all its species regarding their phytochemicals and biological activities, we attempted in this work to summarize and review (1) the phytochemical profile of all plant species belonging to the genus *Galium*, (2) the pharmacological properties of the nutraceuticals isolated from these medicinal plants, (3) the structure-activity relationship between the bioactive compounds’ chemical structure and their mechanism of activity. Accordingly, this review could be of future value regarding the direction of the research on *Galium* species. In addition to the possible discovery of new drugs of various potential activities that serve the pharmaceutical industry.

METHODS

To achieve their work, the authors searched the keywords *Galium*, Rubiaceae, phytochemistry, pharmacology, and biological activity, from 1995 to 2024 from journals and books accessible in databases such as Scopus, Elton B. Stephens CO, ScienceDirect, Embase, Pubmed, and Medline.






RESULTS

Many studies have been successfully conducted on individual *Galium* species, or comparative researches on up to four species were done, to demonstrate the potential of genus *Galium* in treating various diseases. The results collected from all the published articles were reviewed and summarized in our work, and revealed that these plants are rich in phytochemical compounds that are behind their pharmacological activities such as *G. verum* L. named herb Lady’s Bedstraw which is distributed in North Africa, Europe, and Asia, but it can also be found in the northern U.S. and southern Canada. It is a perennial herbaceous plant of the Rubiaceae family rich in phytochemicals such as iridoid glycosides, triterpenes, anthraquinones, phenolic compounds, tannins, essential oils, and saponins. These bioactive compounds provide *G. verum* L a wide application as choleric, diuretic, and for the treatment of epilepsy and gout [14]. The antioxidant potential of this species was demonstrated by the presence of phenolics mainly gallic acid, flavonoids, and chlorophylls [15]. Moreover, many studies have shown its cytotoxic effect [16], and this activity was tested on colon cancer









Table1. *Galium* species’ distribution worldwide.

<i>Galium</i> species	English name	Distribution	Ref.
<i>G. verum</i>	Lady’s Bedstraw	North Africa, Europe, and Asia, but it can also be found in the northern U.S. and southern Canada	[14]
<i>G. mollugo</i>	Hedge bedstraw or false baby’s breath	North Africa, Europe and North America	[17]
<i>G. aparine</i>	Catchweed bedstraw or cleavers	North America and Eurasia	[18]
<i>G. odoratum</i>	Sweet woodruff	Siberia, Northern and Central Europe, Northern Africa and various regions in the north of Iran	[19]
<i>G. rivale</i>	River bedstraw	Europe to Siberia and Palestine	[20]
<i>G. palustre</i>	Marsh-bedstraw	The Mediterranean area and Western Europe, it has become naturalized in North America and other parts of Europe	[21]
<i>G. tinctorium</i>	Stiff marsh bedstraw	Eastern part of North America, from Texas to Labrador and from Minnesota to Florida, plus eastern and central Mexico and the Dominican Republic	[22]
<i>G. boreale</i>	Northern bedstraw	Northern and central Europe, including Britain, south and west to N. Italy and W. Asia.	[23]
<i>G. spurium</i>	Stickwilly or false cleavers	Europe except the north, western Asia, southern tip of Vancouver Island, across the prairie provinces, and in southwest Ontario and southwest Quebec	[24]








Table 2. Chemical structure of the phytochemicals isolated from various *Galium* species [14–16,19,21,25–44].

Phytochemicals isolated from <i>Galium</i> species	Chemical structure	Biological activity
Phenolic compounds (from the aerial parts of the plant species) Rutin (<i>G.verum</i> blossoms) (<i>G. mollugo</i>)		antioxidant, anticancer, antifungal, antibacterial
Quercetin (<i>G.verum</i> , <i>G. mollugo</i> , <i>G. aparine</i>)		
Luteolin (<i>G. mollugo</i> , <i>G. verum</i>)		
Kaempferol (<i>G. mollugo</i> , <i>G. verum</i>)		
Caffeic acid (<i>G. mollugo</i> , <i>G. verum</i> , <i>G. aparine</i> , <i>G. odoratum</i>)		










(Continued)

Phytochemicals isolated from Galium species	Chemical structure	Biological activity
Gallic acid (<i>G. mollugo</i> , <i>G. verum</i> , <i>G. cruciata</i>)		
Chlorogenic acid (<i>G. mollugo</i> , <i>G. verum</i> , <i>G. aparine</i> , <i>G. odoratum</i> , <i>G. album</i>)		
Diosmetin (<i>G. mollugo</i> , <i>G. verum</i>)		
Coumaric acid (<i>G. mollugo</i> , <i>G. verum</i>)		
Ferulic acid (<i>G. mollugo</i> , <i>G. verum</i>)		
Hesperidin (<i>G. mollugo</i> , <i>G. verum</i>)		
Hispidulin (<i>G. mollugo</i> , <i>G. verum</i>)		
Triterpene Saponins (from the aerial parts of the plant species)		
Rivaloside A (<i>G. rivale</i>)		Antifungal, antioxidant, antineoplastic, antiviral, antibacterial

(Continued)

Phytochemicals isolated from Galium species	Chemical structure	Biological activity
Rivaloside B (<i>G. rivale</i>)		
Betulalbuside A (<i>Galium verum</i>)		
Iridoid glycosides (from the aerial parts of the plant species)		
Asperuloside (<i>G. verum blossoms, G. aparine</i>)		Anti-tumorigenic, sedative anti-hypertensive, anti-inflammatory, antibacterial, antiviral, antioxidant, antifungal
Secogalioside (<i>G. mollugo, G. verum</i>)		
Loganin (<i>G. mollugo, G. verum</i>)		
Monotropein (<i>G. mollugo, G. verum, G. odoratum, G. aparine</i>)		
Scandoside (<i>G. mollugo, G. verum, G. odoratum</i>)		

(Continued)

Phytochemicals isolated from Galium species	Chemical structure	Biological activity
Geniposidic acid (<i>G. mollugo</i> , <i>G. verum</i>)		
Anthraquinones (from the roots of the plant species)		
Quinizarin (Roots of <i>Galium.sinaicum</i>)		
Soranjidiol (Roots of <i>G. sinaicum</i>)		
Physcion (<i>G. verum</i>)		
Phytosterols (from the aerial parts of the plant species)		
β -sitosterol (<i>G. mollugo</i> , <i>G. verum</i> , <i>G. odoratum</i> , <i>G. aparine</i>)		Anti-hypertensive, anti-inflammatory, anti-tumorigenic and antioxidant
Campesterol (<i>G. mollugo</i> , <i>G. verum</i> , <i>G. odoratum</i> , <i>G. aparine</i>)		
Essential oils (from the aerial parts of the plant species)		
Germacrene D (<i>G. verum</i>)		Antioxidant, antibacterial, antifungal, skin disorders, effects on renal, hepatobiliary, gastrointestinal and urinary system
Terpinene (<i>G. verum</i>)		
Thymol (<i>G. odoratum</i> , <i>G. aladaghense</i> , <i>G. incanum</i> , and <i>G. dieckii</i>)		

cells (HT29) and proved that *G. verum* L induces apoptosis and raises the level of Google Scholar in the cancer cells [45]. In addition, *G. verum* L has a protective cardiac activity by preserving the heart's diastolic, and systolic function and cardiac contractility since it alleviates the pro-oxidant production and modulates the antioxidant enzyme activity [25]. The highest level of bioactive compounds and the most potent antioxidant potential is available in the blossoms of *G. verum* mainly due to the presence of iridoids (up to 40.8 ± 2.9 mg AE/g), flavonoids (up to 7.3 ± 0.5 mg QE/g), and polyphenols (up to 27.2 ± 1.5 mg GAE/g). Besides the previous phytochemicals, asperuloside identified in *G. verum* is responsible for the diverse pharmacological effects of the plant mainly anti-hypertensive, anti-inflammatory, and anti-tumorigenic [26]. More studies showed the use of this plant as a diuretic, spasmolytic, choleric, and in the treatment of hepatitis [14]. Other results supported the medicinal importance of *Galium mollugo* also named false baby's breath, hedge bedstraw, or white lady's bedstraw, which is distributed in North Africa, Europe, and North America. This species is rich in phytochemicals mainly triterpenes, anthraquinones, phenolics, and iridoid glycosides in addition to other minor compounds such as saponins, tannins, and essential oils. These secondary metabolites make *G. mollugo* a potent antimicrobial mainly against Gram-positive bacteria. Secogalioside is the most important marker of this species and is behind its diverse therapeutic uses as anti-tumorigenic, sedative anti-hypertensive, anti-inflammatory, antibacterial, antiviral, antioxidant, and antifungal [14,17]. Interestingly, the phytochemical investigation of aerial parts of *G. mollugo*, led to the isolation of nine bioactive compounds mainly Z-mollugoside B, mollugoside A, and E-mollugoside B in addition to the following flavonoids including quercetin-3-O-malonyl-glucoside, quercetin-3-O-glucoside, quercetin-3-O-rhamnoside, apigenin-7-O-glucoside, luteolin-7-O-glucoside, and 2 myricetin-3-O-rhamnoside. These nutraceuticals gave the antioxidant potential to *G. mollugo* since some of them showed significant inhibition of Google Scholar production in fibroblasts exposed to AAPH (2,2'-Azobis(2-amidinopropane) dihydrochloride) [46]. Another valuable *Galium* species is *Galium odoratum*, which was recognized for centuries for treating various ailments mainly microbial infections as tea or dry extract, it was also applied topically for external wounds in traditional medicine and as anti-hypoxic and antimicrobial, and also to improve memory and metabolism, digestive and liver problems, stomach complaints, central nervous system problems and disorders, jaundice, and gout treatment [47]. The pharmacological potential is due to the versatile nutraceuticals mainly polyphenolic compounds such as coumarin, asperuloside, rutin, quercetin, monotropein, scandoside, and kaempferol in addition to around 96 compounds in the essential oil mainly thymol (30.6%) and isothymol (22.8%) [27]. *G. odoratum* is used for urinary, and stomach disorders and as a spasmolytic, expectorant, and against liver and bile diseases [48]. The antioxidant potential of essential oils has gained the interest of the medicinal industry and researchers, and many studies have been done to investigate their protective effect among which the study that showed the effect of *G. odoratum* in reducing the damage of DNA caused by oxidation in cultured human lymphocytes [49]. Its antimicrobial, antiviral, and antifungal activities were documented and

proved that its ethanolic extract has an antimicrobial inhibition zone comparable to that of Gentamicin mainly against Gram-positive bacteria. Another proven effect is the locomotor activity decreasing potential in a dose-dependent manner giving this plant species a significant sedative effect acting on the central nervous system without affecting the muscle tone and coordination [50]. Interestingly, *Galium aparine* L. (catchweed bedstraw) which is native to North America and Eurasia, is widely spread among the *Galium* genus, is included in the British Herbal Pharmacopoeia, and is also named «goosegrass» or cleavers. It is traditionally used for treating many skin ailments and burns in tincture form in different antihomotoxic remedies due to the presence of phytochemicals such as iridoids, polyphenols, and flavonoids mainly quercetin 3-O-rhamnoglucoside-7-O-glucoside [18,28]. The antidiabetic and antioxidant properties of *G. aparine* have been demonstrated in several studies on seed extracts [51]. Several reports revealed that methanolic extracts of this species have anti-cancer effects by targeting the cancer cells only without affecting the normal cells as shown in the study done on breast cancer cells and the result was killing the apoptosis-resistant cancer cells by *G. aparine* plant [52]. Moreover, other bioactive substances were isolated from the roots and rhizome of this plant such as coumarins and anthracene compounds mainly alizarin in the roots responsible for the antifungal and antibacterial activity in addition to the isolation of sesquiterpenoids and fatty acids [29]. Other studies confirmed the use of this plant as antimicrobial, antioxidant, and immunomodulatory due to the presence of flavonoids, polyphenols, and hydroxycinnamic derivatives [18]. *Galium* species information collected in Tables 1 and 2 show their distribution, and phytochemical profile in addition to the diversity of their secondary metabolites' chemical structure and their biological activities.

DISCUSSION

The literature review showed that *Galium* species have wide biological activity and this potential is due to the presence of phytochemicals of versatile structure. But despite the successful use of *Galium* plants in traditional medicine for the treatment of various ailments for centuries, the number of studies referring to their pharmacological potential is limited. A comprehensive study of the pharmacological activities of *G. verum* and *Galium mollugo* has been discussed by Bradic *et al.* (2021) to highlight their diverse potential including anti-cancer effects, besides their effects on central nervous system, hepatobiliary, renal, gastrointestinal, and urinary system, in addition to their antioxidant, antifungal, and antibacterial activities [14]. More published articles revealed that the antioxidant potential of *G. verum*, *G. palustre*, *G. cruciata*, and *G. divaricatum* was related to the high level of flavonoids and phenolics [53]. Interestingly, the broad pharmacological potential of *G. odoratum* is due to the high concentrations of coumarin making that species a commercially grown plant to produce coumarins, since this phytochemical is antitumor, bacteriostatic, antipsoriasis, cytotoxic, hepatoprotective, analgesic, antipyretic, and anti-cancerous. Moreover, coumarin is used in the development of many therapeutic drugs used as anticoagulants [50]. Interestingly, Secogalioside being the major component in *G. mollugo* is responsible for the

biological activities of this plant mainly its anti-tumorigenic, sedative, anti-hypertensive, antioxidant, and antimicrobial [17].

The structure-activity relationship studies showed that flavonoids found in most *Galium* plants have a specific mechanism of action based on increasing intracellular mitochondrial ROS, damaging cellular DNA, and this is related to the effect of *G. verum* on human fibroblast cells which is increasing apoptosis induction [16]. However, the chemical structure of flavonoids relies on the number and position of hydroxy groups on the B ring in addition to the position of the catechol b-ring that influences the biological activity mainly the potency of its antioxidant potential [30]. Other authors noticed that polyphenols work as scavengers of free radicals, regulators of nitric oxide, and hence inhibitors of cell proliferation, and these various mechanisms are behind the preventive effects of *Galium* plants including antioxidant, neurodegenerative, cardiovascular, and anti-cancer. For example, the tea of *G. verum* is recommended in the treatment of larynx and tongue cancer and in decoction form, it was shown beneficial in head, neck, breast, and oral cancer. Besides all the previous nutraceuticals in *Galium* plants, *G. mollugo*, *G. verum*, *G. odoratum*, and *G. aparine* being rich in phytosterols have anti-cancer activity mainly against colon cancer because it was reported that phytosterols induce apoptosis by targeting several signaling pathways [34]. Interestingly, phytosterols act as antitumors by boosting cancer immune recognition, affecting the growth of endocrine tumors that are hormonal-dependent, and altering the biosynthesis of sterols. Besides, these phytoconstituents act by direct inhibition of tumor growth and metastasis, slowing the progression of the cell cycle [35]. Accordingly, since studies have shown that the dietary intake of stigmasterol and β -sitosterol resulted in a decrease in ovarian and esophageal cancer risks, respectively [36], it would be interesting to undergo anticancer studies on *G. mollugo*, *G. verum*, *G. odoratum*, and *G. aparine*, being rich in phytosterols. It was demonstrated that the ethanolic extract of *G. verum* rich in phenolic compounds acts by inducing apoptosis and cell death and this mechanism leads to inhibition of human breast cancer cell proliferation [14]. Concerning the immunomodulatory potential of *G. aparine*, a study done to identify the mechanism of action of this plant, showed that *G. aparine* herb contains bioactive fractions made of polysaccharide complex, pectin complex, and polyphenolic complex that stimulate the activity of immunocompetent blood cells and increase the percentage of proliferation of lymphocytes in the lymphocyte blast transformation Google Scholar especially the 96% ethanolic extract of this plant. In addition, these fractions act as direct scavengers of nitrogen oxide (NO), H_2O_2 (hydrogen peroxide), and (2,2-diphenyl-1-picrylhydrazyl) DPPH radical giving the cleavers their antioxidant potential [18].

Other researchers revealed that iridoids exhibit a variety of functionalities including immunomodulatory, antidiabetic, hepatoprotective, choleric, neuroprotective, and antibacterial activities [20]. Concerning asperuloside, an iridoid isolated from some *Galium* species, it was shown to be

responsible for the plant sedative effects, and based on that data, *G. verum* was proven beneficial in nervousness and phobias [14]. Moreover, other researchers investigated the importance of ether terpenes which are responsible for the anti-tumor activity of iridoids because they can inhibit the activity of DNA polymerase [54]. Interestingly, the versatile health benefits of iridoids were shown to be related to the presence of specific oxidative substituents in the cyclopentane ring of the iridoids [55].

More studies were done also to evaluate the anthraquinone mechanism as bacteriostatic and showed that these compounds inhibit bacterial biofilm formation, protein, and nucleic acid synthesis and cause the destruction of bacterial cell walls [56]. These phytochemicals have an anticancer activity which is related to the substitutions of the anthraquinone ring mainly at C-1, C-2, C-3, and C-6 positions, and their anti-pathogenic effect is related to the substitutions at C-1 and C-2 positions while the substitution of the hydroxyl group on the anthraquinone ring is related to most of the activities including the antioxidant and anti-tumor. Moreover, as the polarity of the anthraquinones is stronger, the better is the antibacterial potential of the *Galium* species rich in anthraquinones such as *Galium sinaicum*, *G. verum*, and *G. mollugo* against drug-resistant bacteria including Gram-positive and Gram-negative bacteria. Anthraquinones also exhibit their antibacterial activity by the inhibition of nucleic acids and protein synthesis, and of the respiratory metabolism of bacteria [31]. Furthermore, triterpene saponins have a wide antifungal and antimicrobial potential associated with a specific moiety mainly the aglycone part, and the hydroxyl group esterification affects their effect [32]. These phytoconstituents have a wide biological activity including anti-tumor, anti-inflammatory, and antiviral, and these effects are related to a specific moiety mainly the type of aglycone and the number of sugar chains. The data revealed the antimicrobial properties of some *Galium* species such as *Galium rivale*, are due to the interaction between the sterols on the bacterial erythrocyte membrane and the saponins [33].

Recent studies demonstrated that the antibacterial activity of *Galium* species mainly *Galium aladaghense*, *Galium incanum*, and *Galium dieckii* is related also to the essential oils obtained from the whole plant, this essential oil comprised around 61 compounds including sesquiterpene (14.75%), monoterpenoid (14.75%), monoterpene (8.2%), and these metabolites exhibit a potent inhibitory potential against Gram-negative and Gram-positive bacteria including *Staphylococcus aureus* (*S. aureus*), *Escherichia coli* (*E. coli*), *Pseudomonas syringae* (*P. syringae*), *Salmonella Typhimurium* (*S. typhimurium*), and *Streptococcus mutans* (*S. mutans*) [44]. Accordingly, the mechanism of antibacterial effect is related to a specific moiety, especially the Phenolic-OH present in the essential oil or in the phenolic compounds in addition to the hydrophobicity of the aromatic ring that increases the destruction of the yeast plasma membrane driving to a higher antifungal activity [38]. In conclusion, the structure-activity relationship of the phytochemicals obtained from plants of this genus was demonstrated to illustrate the rationale behind their medicinal uses. The diversity of the

constituents and their evolving biological activities could pave the way for advancing the field of drug discovery and enriching the pharmaceutical industry worldwide. The data illustrated in this study were compiled in a systematic manner to show all the phytochemicals and biological effects collectively. Accordingly, the significance of the analyses done on the collective data facilitates the rapid development of phytochemical research in the direction of drug discovery based on *Galium* plants. Special emphasis on anti-cancer activities to offer a pool of sustainable data that could serve the continuous needs of novel drugs in this regard.

CONCLUSION

Based on the previously mentioned information, the present study demonstrates that *Galium* plants are rich in secondary metabolites of versatile chemical structures including phenolic compounds, triterpene saponins, phytosterols, anthraquinones, and essential oils responsible for the pharmacological potential of these species. The most prominent studies highlighted their antimicrobial, anti-cancer, antioxidant, and anti-inflammatory properties which are all related to a specific moiety in the phytoconstituents such as the phenolic-OH that is common in almost all *Galium* species and which affects the antioxidant and antibacterial potential of these plants based on the number and position of these groups, in addition to the position of substitution of the hydroxyl group on the anthraquinone ring that affects the *Galium* activities mainly those rich in anthraquinones as *G. sinaicum*, *G. verum* and *G. mollugo*. As well as the number of thymol groups, the type of aglycone, and the number of sugar chains in other *Galium* species rich in essential oils and saponins such as *G. odoratum*, *G. aladaghense*, *G. incanum*, *G. dieckii*, and *G. rivale*. Accordingly, based on the diversity of therapeutic uses of *Galium*, and since there is a lack of comprehensive studies that summarized the pharmacological potential of *Galium* species, our work is considered as a cornerstone that opens the way for novel approaches in drug discovery based on *Galium* plants growing worldwide. Accordingly, it is highly recommended that future studies investigate the anti-cancer effect of *Galium* plants to make them possible candidates for anti-cancer medicinal products, especially since few researches have been performed on the effect of this plant on cancer cell lines.

ACKNOWLEDGMENTS

The authors thank Beirut Arab University for providing the facilities during the manuscript preparation. The authors express their gratitude to Biorender.com (free trial version) for enabling the creation of all the figures.

AUTHOR CONTRIBUTIONS

MAK, MAH, AL, MA, made a significant contribution to the work reported, whether that is in the conception, execution, acquisition, analysis, or interpretation of data, or all the areas; took part in drafting, revising, or critically reviewing the article; and gave final approval of the version to be published. All have read and agreed to the published version of the manuscript.

FINANCIAL SUPPORT

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

CONFLICTS OF INTEREST

The authors report no financial or any other conflicts of interest in this work.

ETHICAL APPROVALS

This study does not involve experiments on animals or human subjects.

DATA AVAILABILITY

The data that support the findings of this study are available in standard research databases such as PubMed, Science Direct, or Google Scholar, and/or on public domains that can be searched with either key words or DOI numbers.

USE OF ARTIFICIAL INTELLIGENCE (AI)-ASSISTED TECHNOLOGY

The authors declare that they have not used artificial intelligence (AI)-tools for writing and editing of the manuscript, and no images were manipulated using AI.

PUBLISHER'S NOTE

All claims expressed in this article are solely those of the authors and do not necessarily represent those of the publisher, the editors and the reviewers. This journal remains neutral with regard to jurisdictional claims in published institutional affiliation.

REFERENCES

1. Delprete PG, Jardim JG. Systematics, taxonomy and floristics of Brazilian Rubiaceae: an overview about the current status and future challenges. *Rodriguésia*. 2012;63:101–28. doi: <https://doi.org/10.1590/S2175-78602012000100009>
2. Bolzani VDS, Young MCM, Furlan M, Cavaleiro AJ, Araújo AR, Silva DHS *et al.* Secondary metabolites from Brazilian Rubiaceae plant species: chemotaxonomical and biological significance. *Recent Res Dev Phytochem*. 2001;5:19–31.
3. Heitzman ME, Neto CC, Winiarz E, Vaisberg AJ, Hammond GB. Ethnobotany, phytochemistry and pharmacology of *Uncaria* (Rubiaceae). *Phytochemistry*. 2005 Jan 1;66(1):5–29.
4. Martins D, Nunez CV. Secondary metabolites from Rubiaceae species. *Molecules*. 2015 Jul 22;20(7):13422–95.
5. Sanders JP, Dawson WT. Efficacy of quinidine in malaria. *J Am Med Assoc*. 1932 Nov 19;99(21):1773–7.
6. Aboul-Ela MA, Kamal SA, Badr JM. Growing in Egypt. *Alexandria J Pharm Sci*. 2006 Sep;20(2):79.
7. El-Lakany AM, Aboul-Ela MA, Abdel-Kader MS, Badr JM, Sabri NN, Goher Y. Anthraquinones with antibacterial activities from *Crucianella maritima* L. growing in Egypt. *Nat Prod Sci*. 2004;10(2):63–8.
8. Jansen S, Robbrecht E, Beeckman H, Smets E. A survey of the systematic wood anatomy of the Rubiaceae. *IAWA J*. 2002 Jan 1;23(1):1–67.
9. Judkevich MD, Salas RM, Gonzalez AM. Anatomy and development of the edible fruits of *Cordia* *concolor* (Rubiaceae). *Anais da Academia Brasileira de Ciências*. 2022 Aug 1;94:e20210071.
10. Bremer B, Eriksson T. Time tree of Rubiaceae: phylogeny and dating the family, subfamilies, and tribes. *Int J Plant Sci*. 2009 Jul;170(6):766–93.

11. González-Tejero MR, Casares-Porcel M, Sánchez-Rojas CP, Ramiro-Gutiérrez JM, Molero-Mesa J, Pieroni A, *et al.* Medicinal plants in the Mediterranean area: synthesis of the results of the project Rubia. *J Ethnopharmacol.* 2008 Mar 5;116(2):341–57.
12. Güvenalp Z, Kilic N, Kazaz C, Kaya Y, Demirezer L. Chemical constituents of *Galium tortumense*. *Turkish J Chem.* 2006;30(4):515–23.
13. Ilyina TV, Goryacha OV, Toryanik EL, Kulish IA, Kovaleva AM. Antimicrobial activity of the genus *Galium* L. *Pharm Commun.* 2016;6(1):41–7.
14. Bradic J, Petkovic A, Tomovic M. Phytochemical and Pharmacological properties of some species of the genus. *Exp Appl Biomed Res (EABR).* 2021;22(3):187–93.
15. Lakić NS, Mimica-Dukić NM, Isak JM, Božin BN. Antioxidant properties of *Galium verum* L.(Rubiaceae) extracts. *Central Eur J Bio.* 2010 Jun;5:331–7.
16. Pashapour S, Heshmati M, Mousavi Z, Esmaili S. Effect of whole methanolic extract of *Galium verum* on AGO cell line. *Toxicol Commun.* 2022;4(2):10.
17. Milić PS, Rajković KM, Stamenković OS, Veljković VB. Kinetic modeling and optimization of maceration and ultrasound-extraction of resinoid from the aerial parts of white lady's bedstraw (*Galium mollugo* L.). *Ultrasonics sonochemistry.* 2013 Jan 1;20(1):525–34.
18. Bokhari J, Khan MR, Shabbir M, Rashid U, Jan S, Zai JA. Evaluation of diverse antioxidant activities of *Galium aparine*. *Spectr Acta Part A Mole Biomole Spectr.* 2013 Feb 1;102:24–9.
19. Frišić M, Štibrčić Baglama M, Milović M, Hazler Pilepić K, Maleš Ž. Content of bioactive constituents and antioxidant potential of *Galium* L. species. *Croatica Chemica Acta.* 2018 Dec 29;91(3):411–7.
20. Cooperrider TS. Ohio's herbaria and the Ohio Flora Project. *Ohio J Sci.* 1984;84(4):189–96.
21. Dias MC, Pinto DC, Silva AM. Plant flavonoids: chemical characteristics and biological activity. *Molecules.* 2021 Sep 4;26(17):5377.
22. Dinda B. Pharmacology and applications of naturally occurring iridoids. Berlin/Heidelberg, Germany: Springer International Publishing; 2019 Mar 26.
23. Qun T, Zhou T, Hao J, Wang C, Zhang K, Xu J, *et al.* Antibacterial activities of anthraquinones: structure–activity relationships and action mechanisms. *RSC Med Chem.* 2023;14(8):1446–71.
24. Li Y, Jiang JG. Health functions and structure–activity relationships of natural anthraquinones from plants. *Food Funct.* 2018;9(12):6063–80.
25. Bradic J, Jeremic N, Petkovic A, Jeremic J, Zivkovic V, Srejevic I, *et al.* Cardioprotective effects of *Galium verum* L. extract against myocardial ischemia-reperfusion injury. *Arch Physiol Biochem.* 2020 Oct 19;126(5):408–15.
26. Laanet PR, Saar-Reismaa P, Jõul P, Bragina O, Vaher M. Phytochemical screening and antioxidant activity of selected Estonian *Galium* species. *Molecules.* 2023 Mar 22;28(6):2867.
27. Ilina T, Skowrońska W, Kashpur N, Granica S, Bazylko A, Kovalyova A, *et al.* Immunomodulatory activity and phytochemical profile of infusions from Cleavers herb. *Molecules.* 2020 Aug 14;25(16):3721.
28. Özmatara MB. The effect of extraction methods on antioxidant and enzyme inhibitory activities and phytochemical components of *Galium aparine* L. *Trakya Univ J Nat Sci.* 2021 Apr 15.
29. Kahkeshani N, Farahanikia B, Mahdavian P, Abdolghaffari A, Hassanzadeh G, Abdollahi M, *et al.* Antioxidant and burn healing potential of *Galium odoratum* extracts. *Res Pharm Sci.* 2013 Jul;8(3):197.
30. Ferrer-Gallego PP. Typification of Linnaean specific names in the genus *Galium* (Rubiaceae). *Taxonomy.* 2020 Oct;69(5):1062–71.
31. Mitova MI, Anchev ME, Handjieva NV, Popov SS. Iridoid patterns in *Galium* L. and some phylogenetic considerations. *Zeitschrift für Naturforschung C.* 2002 Apr 1;57(3–4):226–34.
32. El-Gamal AA, Takeya K, Itokawa H, Halim AF, Amer MM, Saad HE, *et al.* Anthraquinones from *Galium sinaicum*. *Phytochemistry.* 1995 Sep 1;40(1):245–51.
33. Biswas T, Dwivedi UN. Plant triterpenoid saponins: biosynthesis, in vitro production, and pharmacological relevance. *Protoplasma.* 2019 Nov;256:1463–86.
34. Wang C, Gong X, Bo A, Zhang L, Zhang M, Zang E, *et al.* Iridoids: research advances in their phytochemistry, biological activities, and pharmacokinetics. *Molecules.* 2020 Jan 10;25(2):287.
35. Elhaw MH, Aldinary MM. Physiological and biochemical studies on *Galium sinaicum* plant. *Al-Azhar Bull Sci.* 2018 Dec 1;29(2-C):63–72.
36. Zengin G, Degirmenci NS, Alpsoy L, Aktumsek A. Evaluation of antioxidant, enzyme inhibition, and cytotoxic activity of three anthraquinones (alizarin, purpurin, and quinizarin). *Human Exp Toxicol.* 2016 May;35(5):544–53.
37. Mocan A, Crisan G, Vlase L, Ivanescu B, Badarau AS, Arsene AL. Phytochemical investigations on four *Galium* species (Rubiaceae) from Romania. *Farmacia.* 2016 Jan 1;64(1):95–9.
38. Karakaş FP, Yildirim A, Türker A. Biological screening of various medicinal plant extracts for antibacterial and antitumor activities. *Turkish journal of biology.* 2012;36(6):641–52.
39. Saha S, Walia S, Kumar J, Parmar BS. Structure–biological activity relationships in triterpenic saponins: the relative activity of protobassic acid and its derivatives against plant pathogenic fungi. *Pest Manag Sci.* 2010 Aug;66(8):825–31.
40. Sparg S, Light ME, Van Staden J. Biological activities and distribution of plant saponins. *J Ethnopharmacol.* 2004 Oct 1;94(2–3):219–43.
41. Woyengo TA, Ramprasath VR, Jones PJ. Anticancer effects of phytosterols. *Eur J Clin Nutr.* 2009 Jul;63(7):813–20.
42. Shahzad N, Khan W, Shadab MD, Ali A, Saluja SS, Sharma S, *et al.* Phytosterols as a natural anticancer agent: current status and future perspective. *Biomed Pharm.* 2017 Apr 1;88:786–94.
43. Yağız F, Battaloğlu R, İlk S, Savran A. Antibacterial activity and chemical composition of essential oils from some *Galium* (Rubiaceae) species against pathogenic bacteria. *Turkish J Agric-Food Sci Technol.* 2017 Oct 30;5(11):1330–3.
44. Konuk HB, Ergüden B. Phenolic–OH group is crucial for the antifungal activity of terpenoids via disruption of cell membrane integrity. *Folia Microbio.* 2020 Aug;65(4):775–83.
45. Pashapour S, Heshmati M, Mousavi Z, Esmaili S. The apoptotic effect of methanolic extract of *Galium verum* on HT29 cell line. *J Biol Stud.* 2022 Mar 30;4(4):210–20.
46. Chaher N, Krisa S, Delaunay JC, Bernillon S, Pedrot E, Mérillon JM, *et al.* Unusual compounds from *Galium mollugo* and their inhibitory activities against ROS generation in human fibroblasts. *J Pharm Biomed Anal.* 2016 Jan 5;117:79–84.
47. Bavya S, Nagalakshmi R, Sundari UT. Anatomical and preliminary phytochemical evaluation of the medicinal plant: *enicostemma littorale*. *Medicinal.* 2022;31.
48. Marković MS, Pljevljakušić DS, Pančić AS, Rakonjac LB, Nikolić BM, Stankov JV. Ethnobotanical use of plants from the genus *Galium* in the Pirot District. *Pirotski Zbornik.* 2023(48):191–202.
49. Khabour OF, Alzoubi KH, Hassanein SF, Makhoulouf H, Alhashimi F. Protective effect of essential oils of *ocimum basilicum*, *Galium odoratum*, and *cymbopogon citratus* against oxidative dna damage in cultured human lymphocyte cells. *Pak J Bot.* 2023 Mar 3;55(1):307–11. doi: [http://dx.doi.org/10.30848/PJB2023-1\(4\)](http://dx.doi.org/10.30848/PJB2023-1(4))
50. Ilina T, Kashpur N, Granica S, Bazylko A, Shinkovenko I, Kovalyova A, *et al.* Phytochemical profiles and in vitro immunomodulatory activity of ethanolic extracts from *Galium aparine* L. *Plants.* 2019 Nov 25;8(12):541.
51. Atmaca H, Bozkurt E, Citan M, Tepe HD. Effects of *Galium aparine* extract on the cell viability, cell cycle and cell death in breast cancer cell lines. *J Ethnopharmacol.* 2016 Jun 20;186:305–10.
52. Goryacha OV, Ilyina TV, Kovalyova AM, Kashpur NV. Phytochemical research of *Galium aparine* L. lipophilic complex and study of its antibacterial activity. *Pharma Innov.* 2014;3:1.

53. De Rosa S, Iodice C, Mitova M, Handjieva N, Popov S, Anchev M. Triterpene saponins and iridoid glucosides from *Galium rivale*. *Phytochemistry*. 2000 Aug 1;54(8):751–6.
54. Wood S, Gould K, Smreciu A. *Galium boreale*: northern bedstraw, sweet scented bedstraw, fragrant bedstraw. Edmonton, Canada, University of Alberta Library 2013.
55. Orhan N, Orhan DD, Aslan M, Şüküroğlu M, Orhan IE. UPLC–TOF-MS analysis of *Galium spurium* towards its neuroprotective and anticonvulsant activities. *J Ethnopharmacol*. 2012 May 7;141(1):220–7.
56. De Rosa S, Mitova M, Handjieva N, Popov S, Anchev M. Rivalosides A and B, two 19-oxo triterpenoid saponins from *Galium rivale*. *J Nat Prod*. 2000 Jul 28;63(7):1012–4.

How to cite this article:

Kanso MA, Hijazi MA, El-Lakany A, Aboul-Ela M. Review on phytochemical constituents and pharmacological activities of genus *Galium*. *J Appl Pharm Sci*. 2024;14(09):046–056.